

# Exhibit A



US 20100130591A1


(19) **United States**(12) **Patent Application Publication**  
**Sazani et al.**(10) **Pub. No.: US 2010/0130591 A1**(43) **Pub. Date: May 27, 2010**(54) **MULTIPLE EXON SKIPPING  
COMPOSITIONS FOR DMD****Publication Classification**(76) Inventors: **Peter Sazani**, Bothell, WA (US);  
**Ryszard Kole**, Bellevue, WA (US)(51) **Int. Cl.****A61K 31/7088** (2006.01)**C07H 21/02** (2006.01)**C07K 9/00** (2006.01)**A61P 21/00** (2006.01)

Correspondence Address:


**SEED INTELLECTUAL PROPERTY LAW  
GROUP PLLC****701 FIFTH AVE, SUITE 5400****SEATTLE, WA 98104 (US)**(52) **U.S. Cl. .... 514/44 A; 536/24.5; 530/322**(21) Appl. No.: **12/605,276**(22) Filed: **Oct. 23, 2009**

(57)

**ABSTRACT****Related U.S. Application Data**(60) Provisional application No. 61/108,416, filed on Oct.  
24, 2008.Provided are antisense molecules capable of binding to a  
selected target site in the human dystrophin gene to induce  
exon skipping, and methods of use thereof to treat muscular  
dystrophy.



*FIG. 1A*



**FIG. 1B**







FIG. 1C




*FIG. 1D*



*FIG. 1E*



*FIG. 1F*



*FIG. 1G*

# Dystrophin Exon 51 Scan Oligos

DMD Gene 1756



Oligo SEQ ID NOs shown – SEQ ID NOs 324, 326 and 327 were most effective

*FIG. 2A*

# Exon 51S.3 (RD)

High-Purity Synthesis, 3.0uM, RD Cells

Oligo	Name; SEQ ID NO	Lot
NG-07-1160	AVI-5658; 588	09MY11-R(E4)
NG-09-0053	053; 324	09JNJ12-R(A4)
NG-09-0054	054; 326	09JNJ12-R(B4)
NG-09-0055	055; 327	09JNJ12-R(E4)

NG-09-0053 (SEQ ID NO:324)	NG-09-0054 (SEQ ID NO:326)	NG-09-0055 (SEQ ID NO:327)	NG-07-1160 (SEQ ID NO:588)
4.65% ±1.89	7.40% ±0.75	9.89% ±1.37	5.26% ±0.66




FIG. 2B



51MCS (Muscle Cell Competitor Screen)  
High-Purity Synthesis, 3.0uM, Human Primary Skeletal Muscle Cells

Oligo	Name; SEQ ID NO	Lot
NG-07-1160	AVI-5658; 588	09MY11-R(E4)
NG-09-0053	053; 324	09JNJ12-R(A4)
NG-09-0054	054; 326	09JNJ12-R(B4)
NG-09-0055	055; 327	09JNJ12-R(E4)
NG-08-0835	h51AON1; 594	09JL07-R(A1)




FIG. 2C

# Selected Dystrophin Exon 51 Oligos

DMD Gene 1756




FIG. 2D

# Dystrophin Exon 50 Scan Oligos

DMD Gene 1756




FIG. 3A

Compound	EC <sub>50</sub> (micromolar)*
AVI-5656 (SEQ ID NO:584)	0.921
AVI-5915 (SEQ ID NO:585)	3.693
NG-08-0731 (SEQ ID NO:277)	1.741
AVI-5038 (SEQ ID NO:287)	0.966
NG-08-0741 (SEQ ID NO:290)	1.836
NG-08-0742 (SEQ ID NO:291)	2.402

\*Determined from dose-ranging studies in RD cells  
AVI #00453 - 02FEB2009

FIG. 3B

## Dystrophin Exon 53 Scan Oligos

DMD Gene 1756




FIG. 4A

Selected Dystrophin Exon 53 Oligos




FIG. 4B

Exon 53 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0746 09AU11-R(E5) (SEQ ID NO: 422)



FIG. 4C

Exon 53 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0749 09AU11-R(A7) (SEQ ID NO: 428)




FIG. 4D



Exon 53 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0750 09AU11-R(C7) (SEQ ID NO: 429)




FIG. 4E

Exon 53 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0751 09AU11-J(E7) (SEQ ID NO: 431)



FIG. 4F

Exon53 RD Cell Dose-Range Summary

Percent Exon Skipped					
Treatment (uM)	746 (SEQ ID NO:422)	749 (SEQ ID NO:428)	750 (SEQ ID NO:429)	751 (SEQ ID NO:431)	
1.0	0.00	1.05	4.83	0.79	
2.0	0.00	1.72	10.44	1.98	
3.0	0.00	2.40	17.28	2.74	
5.0	0.54	2.62	23.29	4.18	
10	3.53	4.63	57.67	13.89	
EC <sub>50</sub> (uM)	NA	72.2	9.3	25.3	

FIG. 4G

Exon 53 CS (Competitor Screen)  
High-Purity Synthesis, 3.0uM, RD Cells




FIG. 4H

Exon 53 MCS (Muscle Cell Competitor Screen)  
High-Purity Synthesis, 3.0uM, Human Primary Skeletal Muscle Cells

Oligo	Name; SEQ ID	Lot
NG-09-0490	H53A(+39+69); 608	09AU18-J(A1)
NG-08-0750 (W)	H53A(+23+47); 609	09-AU18-J(B1)
NG-09-0559	h53AON1; 610	09AU18-J(C1)
NG-09-0560	H53A(-12+10); 611	09AU18-J(D1)
NG-08-0750	008; 429	09AU11-R(C7)




FIG. 4I

## Dystrophin Exon 44 Scan Oligos




FIG. 5A

# Selected Dystrophin Exon 44 Oligos





FIG. 5B

**Exon 44 DR.2 (Dose-Range)**  
High-Purity Synthesis, (0.1, 0.3, 1.0, 3.0, 10.0uM) RD Cells

NG-08-0792 09AU11-J(A2) (SEQ ID NO: 4)



*FIG. 5C*



Exon 44 DR.2

High-Purity Synthesis, (0.1, 0.3, 1.0, 3.0, 10.0uM) RD Cells

NG-08-0796 09AU11-J(B2) (SEQ ID NO: 8)




FIG. 5D

Exon 44 DR.2  
High-Purity Synthesis, (0.1, 0.3, 1.0, 3.0, 10.0uM) RD Cells  
NG-08-0799 09AU11-J(C2) (SEQ ID NO: 11)




FIG. 5E

# Exon 44 DR.2

High-Purity Synthesis, (0.1, 0.3, 1.0, 3.0, 10.0uM) RD Cells

NG-08-0800 09AU11-J(D2) (SEQ ID NO: 12)




FIG. 5F

Exon 44 DR.2

High-Purity Synthesis, (0.1, 0.3, 1.0, 3.0, 10.0uM) RD Cells

NG-0801 09AU11-J(E2) (SEQ ID NO: 13)




FIG. 5G

# Exon44 RD Cell Dose-Range Summary

Percent Exon Skipped

Treatment (uM)	792 (SEQ ID NO:4)	796 (SEQ ID NO:8)	799 (SEQ ID NO:11)	800 (SEQ ID NO:12)	801 (SEQ ID NO:13)
0.1	3.14	4.14	3.00	7.49	2.10
0.3	4.66	6.69	8.41	16.01	4.06
1	7.56	17.41	21.48	18.81	9.01
3	21.64	39.74	32.92	46.71	22.50
10	64.94	97.39	88.47	96.52	69.38
EC <sub>50</sub> (uM)	7.166	3.024	3.594	2.795	6.277

FIG. 5H

Exon 44 CS (Competitor Screen)

High-Purity Synthesis, 3.0uM, RD Cells

Oligo	Name; Seq ID	Lot
NG-09-0561	H44A(+61+84); 600	09AU18-J(E1)
NG-09-0562	H44A(+85+104); 601	09AU18-J(F1)
NG-09-0563	h44AON1; 602	09AU18-J(A2)
NG-09-0564	H44A(-06+14); 603	09AU18-J(B2)
NG-08-0800	012; 12	09AU11-J(D2)

561	562	563	564	800
SEQ ID NO: 600	SEQ ID NO: 601	SEQ ID NO: 602	SEQ ID NO: 603	SEQ ID NO: 12
44.4% ±4.22	20.4% ±0.84	94.0% ±0.32	97.1% ±0.07	47.5% ±2.05
Not quantified				
MW 0				




FIG. 5I

Exon 44 MCS (Muscle Cell Competitor Screen)  
High-Purity Synthesis, 3.0uM, Human Primary Skeletal Muscle Cells

Oligo	Name; Seq ID	Lot
NG-09-0561	H44A(+61+84); 600	09AU18-J(E1)
NG-09-0562	H44A(+85+104); 601	09AU18-J(F1)
NG-09-0563	h44AON1; 602	09AU18-J(A2)
NG-09-0564	H44A(-06+14); 603	09AU18-J(B2)
NG-08-0800	012; 12	09AU11-J(D2)

Inadequate NG-09-0564 to include in screen.




FIG. 5J

## Dystrophin Exon 45 Scan Oligos

DMD Gene 1756





FIG. 6A



## Selected Dystrophin Exon 45 Oligos



*FIG. 6B*

# Exon 45 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0770 09AU11-J(F2) (SEQ ID NO: 27)




FIG. 6C

Exon 45 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0771 09AU11-J(A4) (SEQ ID NO: 29)




FIG. 6D

Exon 45 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0774 09AU11-J(B4) (SEQ ID NO: 34)




FIG. 6E

# Exon 45 DR.2

High-Purity Synthesis, (1.0, 2.0, 3.0, 5.0, 10.0uM) RD Cells

NG-0777 09AU11-J(C4) (SEQ ID NO: 39)




FIG. 6F




FIG. 6G

# Exon45 RD Cell Dose-Range Summary

Percent Exon Skipped						
Treatment (uM)	770 (SEQ ID NO:27)	771 (SEQ ID NO:29)	774 (SEQ ID NO:34)	777 (SEQ ID NO:39)	782 (SEQ ID NO:49)	
1.0	0.00	0.53	3.05	0.00	0.00	
2.0	0.43	1.99	13.33	0.00	0.00	
3.0	0.60	4.91	31.12	0.55	0.00	
5.0	1.15	14.25	79.20	1.63	0.78	
10	4.17	84.37	97.68	7.86	2.11	
EC <sub>50</sub> (uM)	54.37	7.25	3.69	NA	NA	

*FIG. 6H*





Exon 45 MCS (Muscle Cell Competitor Screen)  
High-Purity Synthesis, 3.0uM, Human Primary Skeletal Muscle Cells

Oligo	Name; SEQ ID	Lot
NG-09-0565	H45A(+71+90); 604	09AU18-J(C2)
NG-09-0566	h45AON1; 605	09AU18-J(D2)
NG-09-0567	h45AON5; 606	09AU18-J(E2)
NG-09-0568	H45A(-06+20); 607	09AU18-J(F2)
NG-08-0774	008; 34	09AU11-J(B4)




FIG. 6J

US 2010/0130591 A1

May 27, 2010

1

# **MULTIPLE EXON SKIPPING COMPOSITIONS FOR DMD**

## **CROSS-REFERENCE TO RELATED APPLICATION**

**[0001]** This application claims the benefit under 35 U.S.C. §119(e) of U.S. Provisional Patent Application No. 61/108,416 filed Oct. 24, 2008; wherein this provisional application is incorporated herein by reference in its entirety.

## **STATEMENT REGARDING SEQUENCE LISTING**

**[0002]** The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is 120178\_410\_SEQUENCE\_LISTING.txt. The text file is 157 KB, was created on Oct. 23, 2009 and is being submitted electronically via EFS-Web.

## **FIELD OF THE INVENTION**

**[0003]** The present invention relates to novel antisense compounds and compositions suitable for facilitating exon skipping in the human dystrophin gene. It also provides methods for inducing exon skipping using the antisense compositions adapted for use in the methods of the invention.

## **BACKGROUND OF THE INVENTION**

**[0004]** Antisense technologies are being developed using a range of chemistries to affect gene expression at a variety of different levels (transcription, splicing, stability, translation). Much of that research has focused on the use of antisense compounds to correct or compensate for abnormal or disease-associated genes in a wide range of indications. Antisense molecules are able to inhibit gene expression with specificity, and because of this, many research efforts concerning oligonucleotides as modulators of gene expression have focused on inhibiting the expression of targeted genes or the function of cis-acting elements. The antisense oligonucleotides are typically directed against RNA, either the sense strand (e.g., mRNA) or minus-strand in the case of some viral RNA targets. To achieve a desired effect of specific gene down-regulation, the oligonucleotides generally either promote the decay of the targeted mRNA, block translation of the mRNA or block the function of cis-acting RNA elements thereby effectively preventing either de novo synthesis of the target protein or replication of the viral RNA.

**[0005]** However, such techniques are not useful where the object is to up-regulate production of the native protein or compensate for mutations that induce premature termination of translation such as nonsense or frame-shifting mutations. In these cases, the defective gene transcript should not be subjected to targeted degradation or steric inhibition, so the antisense oligonucleotide chemistry should not promote target mRNA decay or block translation.

**[0006]** In a variety of genetic diseases, the effects of mutations on the eventual expression of a gene can be modulated through a process of targeted exon skipping during the splicing process. The splicing process is directed by complex multi-component machinery that brings adjacent exon-intron junctions in pre-mRNA into close proximity and performs cleavage of phosphodiester bonds at the ends of the introns with their subsequent reformation between exons that are to

be spliced together. This complex and highly precise process is mediated by sequence motifs in the pre-mRNA that are relatively short semi-conserved RNA segments to which bind the various nuclear splicing factors that are then involved in the splicing reactions. By changing the way the splicing machinery reads or recognizes the motifs involved in pre-mRNA processing, it is possible to create differentially spliced mRNA molecules. It has now been recognized that the majority of human genes are alternatively spliced during normal gene expression, although the mechanisms involved have not been identified.

**[0007]** In cases where a normally functional protein is prematurely terminated because of mutations therein, a means for restoring some functional protein production through antisense technology has been shown to be possible through intervention during the splicing processes, and that if exons associated with disease-causing mutations can be specifically deleted from some genes, a shortened protein product can sometimes be produced that has similar biological properties of the native protein or has sufficient biological activity to ameliorate the disease caused by mutations associated with the exon (Sierakowska, Sambade et al. 1996; Wilton, Lloyd et al. 1999; van Deutekom, Bremmer-Bout et al. 2001; Lu, Mann et al. 2003; Aartsma-Rus, Janson et al. 2004). Kole et al. (U.S. Pat. Nos. 5,627,274; 5,916,808; 5,976,879; and 5,665,593) disclose methods of combating aberrant splicing using modified antisense oligonucleotide analogs that do not promote decay of the targeted pre-mRNA. Bennett et al (U.S. Pat. No. 6,210,892) describe antisense modulation of wild-type cellular mRNA processing also using antisense oligonucleotide analogs that do not induce RNase H-mediated cleavage of the target RNA.

**[0008]** The process of targeted exon skipping is likely to be particularly useful in long genes where there are many exons and introns, where there is redundancy in the genetic constitution of the exons or where a protein is able to function without one or more particular exons. Efforts to redirect gene processing for the treatment of genetic diseases associated with truncations caused by mutations in various genes have focused on the use of antisense oligonucleotides that either: (1) fully or partially overlap with the elements involved in the splicing process; or (2) bind to the pre-mRNA at a position sufficiently close to the element to disrupt the binding and function of the splicing factors that would normally mediate a particular splicing reaction which occurs at that element.

**[0009]** Duchenne muscular dystrophy (DMD) is caused by a defect in the expression of the protein dystrophin. The gene encoding the protein contains 79 exons spread out over more than 2 million nucleotides of DNA. Any exonic mutation that changes the reading frame of the exon, or introduces a stop codon, or is characterized by removal of an entire out of frame exon or exons or duplications of one or more exons has the potential to disrupt production of functional dystrophin, resulting in DMD.

**[0010]** A less severe form of muscular dystrophy, Becker muscular dystrophy (BMD) has been found to arise where a mutation, typically a deletion of one or more exons, results in a correct reading frame along the entire dystrophin transcript, such that translation of mRNA into protein is not prematurely terminated. If the joining of the upstream and downstream exons in the processing of a mutated dystrophin pre-mRNA maintains the correct reading frame of the gene, the result is an mRNA coding for a protein with a short internal deletion that retains some activity resulting in a Becker phenotype.

US 2010/0130591 A1

May 27, 2010

2

**[0011]** Deletions of an exon or exons which do not alter the reading frame of a dystrophin protein give rise to a BMD phenotype, whereas an exon deletion that causes a frame-shift will give rise to DMD (Monaco, Bertelson et al. 1988). In general, dystrophin mutations including point mutations and exon deletions that change the reading frame and thus interrupt proper protein translation result in DMD. It should also be noted that some BMD and DMD patients have exon deletions covering multiple exons.

**[0012]** Although antisense molecules may provide a tool in the treatment of Duchenne Muscular Dystrophy (DMD), attempts to induce exon skipping using antisense molecules have had mixed success. Successful skipping of dystrophin exon 19 from the dystrophin pre-mRNA was achieved using a variety of antisense molecules directed at the flanking splice sites or motifs within the exon involved in exon definition as described by Errington et al., (Errington, Mann et al. 2003).

**[0013]** The first example of specific and reproducible exon skipping in the mdx mouse model was reported by Wilton et al (Wilton, Lloyd et al. 1999). By directing an antisense molecule to the donor splice site, exon 23 skipping was induced in the dystrophin mRNA within 6 hours of treatment of the cultured cells. Wilton et al also describe targeting the acceptor region of the mouse dystrophin pre-mRNA with longer antisense oligonucleotides. While the first antisense oligonucleotide directed at the intron 23 donor splice site induced exon skipping in primary cultured myoblasts, this compound was found to be much less efficient in immortalized cell cultures expressing higher levels of dystrophin.

**[0014]** Despite these efforts, there remains a need for improved antisense oligomers targeted to multiple dystrophin exons and improved muscle delivery compositions and methods for DMD therapeutic applications.

#### BRIEF SUMMARY OF THE INVENTION

**[0015]** Embodiments of the present invention relate generally to antisense compounds capable of binding to a selected target to induce exon skipping, and methods of use thereof to induce exon skipping. In certain embodiments, it is possible to combine two or more antisense oligonucleotides of the present invention together to induce single or multiple exon skipping.


**[0016]** In certain embodiments, it is possible to improve exon skipping of a single or multiple exons by covalently linking together two or more antisense oligonucleotide molecules (see, e.g., Aartsma-Rus, Janson et al. 2004).

**[0017]** In certain embodiments, the antisense compounds of the present invention induce exon skipping in the human dystrophin gene, and thereby allow muscle cells to produce a functional dystrophin protein.

**[0018]** The antisense oligonucleotide compounds (also referred to herein as oligomers) of the present invention typically: (i) comprise morpholino subunits and phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, (ii) contain between 10-40 nucleotide bases, preferably 20-35 bases (iii) comprise a base sequence effective to hybridize to at least 12 contiguous bases of a target sequence in dystrophin pre-mRNA and induce exon skipping.

**[0019]** In certain embodiments, the antisense compounds of the present invention may comprise phosphorus-containing intersubunit linkages joining a morpholino nitrogen of

one subunit to a 5' exocyclic carbon of an adjacent subunit, in accordance with the following structure (I):



wherein:

**[0020]** Y<sub>1</sub> is —O—, —S—, —NH—, or —CH<sub>2</sub>—;

**[0021]** Z is O or S;

**[0022]** P<sub>j</sub> is a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide; and

**[0023]** X is fluoro, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted thioalkoxy, amino, optionally substituted alkylamino, or optionally substituted heterocyclyl.

**[0024]** In certain embodiments, the above intersubunit linkages, which are uncharged, may be interspersed with linkages that are positively charged at physiological pH, where the total number of positively charged linkages is between 2 and no more than half of the total number of linkages. For example, the positively charged linkages may have the above structure in which X is optionally substituted 1-piperazinyl. In other embodiments, the positively charged linkages may have the above structure in which X is substituted 1-piperazinyl, wherein the 1-piperazinyl is substituted at the 4-position with an optionally substituted alkyl guanidynyl moiety.

**[0025]** Where the antisense compound administered is effective to target a splice site of preprocessed human dystrophin, it may have a base sequence complementary to a target region containing at least 12 contiguous bases in a preprocessed messenger RNA (mRNA) human dystrophin transcript. Exemplary antisense sequences include those identified by SEQ ID NOS: 1 to 569 and 612 to 633.

**[0026]** In certain embodiments, an antisense sequence of the present invention is contained within:

**[0027]** (a) any of the sequences identified by SEQ ID NOS: 1-20, preferably SEQ ID NOS: 4, 8, 11 and 12, and more preferably SEQ ID NO: 12 for use in producing skipping of exon 44 in the processing of human dystrophin pre-processed mRNA;

**[0028]** (b) any of the sequences identified by SEQ ID NOS: 21-76 and 612 to 624, preferably SEQ ID NOS: 27, 29, 34 and 39, and more preferably SEQ ID NO: 34 for use in producing skipping of exon 45 in the processing of human dystrophin pre-processed mRNA;

**[0029]** (c) any of the sequences identified by SEQ ID NOS: 77-125, preferably SEQ ID NOS: 21 to 53, and more preferably SEQ ID NOS: 82, 84-87, 90 96, 98, 99 and 101, for use in producing skipping of exon 46 in the processing of human dystrophin pre-processed mRNA;

**[0030]** (d) any of the sequences identified by SEQ ID NOS: 126-169, preferably SEQ ID NOS: 126-149, and more preferably SEQ ID NOS: 126, 128-130, 132, 144 and 146-149, for use in producing skipping of exon 47 in the processing of human dystrophin pre-processed mRNA;

US 2010/0130591 A1

May 27, 2010

3

**[0031]** (e) any of the sequences identified by SEQ ID NOS: 170-224 and 634, preferably SEQ ID NOS: 170-201 and 634, and more preferably SEQ ID NOS: 176, 178, 181-183, 194 and 198-201, for use in producing skipping of exon 48 in the processing of human dystrophin pre-processed mRNA;

**[0032]** (f) any of the sequences identified by SEQ ID NOS: 225-266, preferably SEQ ID NOS: 225-248, and more preferably SEQ ID NOS: 227, 229, 234, 236, 237 and 244-248, for use in producing skipping of exon 49 in the processing of human dystrophin pre-processed mRNA;

**[0033]** (g) any of the sequences identified by SEQ ID NOS: 267-308, preferably SEQ ID NOS: 277, 287 and 290, and more preferably SEQ ID NO: 287, for use in producing skipping of exon 50 in the processing of human dystrophin pre-processed mRNA;

**[0034]** (h) any of the sequences identified by SEQ ID NOS: 309-371, preferably SEQ ID NOS: 324, 326 and 327, and more preferably SEQ ID NO: 327 for use in producing skipping of exon 51 in the processing of human dystrophin pre-processed mRNA;

**[0035]** (i) any of the sequences identified by SEQ ID NOS: 372-415, preferably SEQ ID NOS: 372-397, and more preferably SEQ ID NOS: 379-382, 384, 390 and 392-395 for use in producing skipping of exon 52 in the processing of human dystrophin pre-processed mRNA;

**[0036]** (j) any of the sequences identified by SEQ ID NOS: 416-475 and 625-633, preferably SEQ ID NOS: 428, 429 and 431, and more preferably SEQ ID NO: 429, for use in producing skipping of exon 53 in the processing of human dystrophin pre-processed mRNA;

**[0037]** (k) any of the sequences identified by SEQ ID NOS: 476-519, preferably SEQ ID NOS: 476-499, and more preferably SEQ ID NOS: 479-482, 484, 489 and 491-493, for use in producing skipping of exon 54 in the processing of human dystrophin pre-processed mRNA; and

**[0038]** (l) any of the sequences identified by SEQ ID NOS: 520-569 and 635, preferably SEQ ID NOS: 520-546 and 635, and more preferably SEQ ID NOS: 524-528, 537, 539, 540, 542 and 544, for use in producing skipping of exon 55 in the processing of human dystrophin pre-processed mRNA;

**[0039]** In certain embodiments, the compound may be conjugated to an arginine-rich polypeptide effective to promote uptake of the compound into cells. Exemplary peptides include those identified by SEQ ID NOS: 570 to 578, among others described herein.

**[0040]** In one exemplary embodiment, the arginine-rich polypeptide is covalently coupled at its N-terminal or C-terminal residue to the 3' or 5' end of the antisense compound. Also in an exemplary embodiment, the antisense compound is composed of morpholino subunits and phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit.

**[0041]** In general, the peptide-oligomer conjugate may further comprise a homing peptide which is selective for a selected mammalian tissue, i.e., the same tissue being targeted by the cell-penetrating peptide. The conjugate may be of the form: cell penetrating peptide-homing peptide-antisense oligomer, or, more preferably, of the form: homing peptide-cell penetrating peptide-antisense oligomer. For example, a peptide conjugate compound for use in treating Duchenne muscular dystrophy, as described above, can further comprise a homing peptide which is selective for muscle tissue, such as the peptide having the sequence identified as SEQ ID NO: 579, conjugated to the cell-penetrating peptide.

Exemplary conjugates of this type include those represented herein as CP06062-MSP-PMO (cell penetrating peptide-homing peptide-antisense oligomer) and as MSP-CP06062-PMO (homing peptide-cell penetrating peptide-antisense oligomer) (see SEQ ID NOS: 580-583).

**[0042]** In some embodiments, the peptide is conjugated to the oligomer via a linker moiety. In certain embodiments the linker moiety may comprise an optionally substituted piperazinyl moiety. In other embodiments, the linker moiety may further comprise a beta alanine and/or a 6-aminohexanoic acid subunit. In yet other embodiments, the peptide is conjugated directly to the oligomer without a linker moiety.

**[0043]** Conjugation of the peptide to the oligomer may be at any position suitable for forming a covalent bond between the peptide and the oligomer or between the linker moiety and the oligomer. For example, in some embodiments conjugation of the peptide may be at the 3' end of the oligomer. In other embodiments, conjugation of the peptide to the oligomer may be at the 5' end of the oligomer. In yet other embodiments, the peptide may be conjugated to the oligomer through any of the intersubunit linkages.

**[0044]** In some embodiments, the peptide is conjugated to the oligomer at the 5' end of the oligomer. In embodiments comprising phosphorus-containing intersubunit linkages, the peptide may be conjugated to the oligomer via a covalent bond to the phosphorous of the terminal linkage group. Conjugation in this manner may be with or without the linker moiety described above.

**[0045]** In yet other embodiments, the peptide may be conjugated to the oligomer at the 3' end of the oligomer. In some further embodiments, the peptide may be conjugated to the nitrogen atom of the 3' terminal morpholino group of the oligomer. In this respect, the peptide may be conjugated to the oligomer directly or via the linker moiety described above.

**[0046]** In some embodiments, the oligomer may be conjugated to a moiety that enhances the solubility of the oligomer in aqueous medium. In some embodiments, the moiety that enhances solubility of the oligomer in aqueous medium is a polyethyleneglycol. In yet further embodiments, the moiety that enhances solubility of the oligomer in aqueous medium is triethylene glycol. For example, in some embodiments the moiety that enhances solubility in aqueous medium may be conjugated to the oligomer at the 5' end of the oligomer. Conjugation of the moiety that enhances solubility of the oligomer in aqueous medium to the oligomer may be either directly or through the linker moiety described above.

**[0047]** Certain embodiments of the present invention provide antisense molecules selected and or adapted to aid in the prophylactic or therapeutic treatment of a genetic disorder comprising at least an antisense molecule in a form suitable for delivery to a patient.

**[0048]** Certain embodiments of the invention provide methods for treating a patient suffering from a genetic disease wherein there is a mutation in a gene encoding a particular protein and the affect of the mutation can be abrogated by exon skipping, comprising the steps of: (a) selecting an antisense molecule in accordance with the methods described herein; and (b) administering the molecule to a patient in need of such treatment. The present invention also includes the use of purified and isolated antisense oligonucleotides of the invention, for the manufacture of a medicament for treatment of a genetic disease.

**[0049]** Certain embodiments provide a method of treating muscular dystrophy, such as a condition characterized by



US 2010/0130591 A1

May 27, 2010

4

Duchenne muscular dystrophy, which method comprises administering to a patient in need of treatment an effective amount of an appropriately designed antisense oligonucleotide, as described herein, relevant to the particular genetic lesion in that patient. Further, certain embodiments provide a method for prophylactically treating a patient to prevent or at least minimize muscular dystrophy, including Duchenne muscular dystrophy, comprising the step of: administering to the patient an effective amount of an antisense oligonucleotide or a pharmaceutical composition comprising one or more of these biological molecules.

**[0050]** Certain embodiments relate to methods of treating muscular dystrophy in a subject, comprising administering to the subject an effective amount of a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 1 to 569 and 612 to 635, and capable of forming with the complementary mRNA sequence in a dystrophin-gene exon a heteroduplex structure between said compound and mRNA having a Tm of at least 45° C., wherein the exon is selected from the group consisting of exons 44-55.

**[0051]** In certain embodiments, the muscular dystrophy is Duchenne's muscular dystrophy (DMD). In certain embodiments, the muscular dystrophy is Becker muscular dystrophy (BMD).

**[0052]** In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 1-20, and the exon is exon 44. In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 21-76 and 612 to 624, and the exon is exon 45.

**[0053]** In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 77-125, and the exon is exon 46. In certain embodiments, the sequence selected from the group consisting SEQ ID NOS: 126-169, and the exon is exon 47.

**[0054]** In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 170-224 and 634, and the exon is exon 48. In certain embodiments, the sequence selected from the group consisting SEQ ID NOS: 225-266, and the exon is exon 49.

**[0055]** In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 267-308, and the exon is exon 50. In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 309-371, and the exon is exon 51.

**[0056]** In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 372-415, and the exon is exon 52. In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 416-475 and 625-633, and the exon is exon 53. In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 476-519, and the exon is exon 54. In certain embodiments, the sequence is selected from the group consisting SEQ ID NOS: 520-569 and 635, and the exon is exon 55. In certain embodiments, the sequence comprises or consists essentially of SEQ ID NO: 287.

**[0057]** Certain embodiments provide kits for treating a genetic disease, which kits comprise at least an antisense oligonucleotide of the present invention, packaged in a suitable container and instructions for its use.

**[0058]** These and other objects and features will be more fully understood when the following detailed description of the invention is read in conjunction with the figures.

#### BRIEF DESCRIPTION OF THE FIGURES

**[0059]** FIG. 1A shows an exemplary morpholino oligomer structure with a phosphorodiamidate linkage;

**[0060]** FIG. 1B shows a conjugate of an arginine-rich peptide and an antisense oligomer, in accordance with an embodiment of the invention;

**[0061]** FIG. 1C shows a conjugate as in FIG. 1B, wherein the backbone linkages contain one or more positively charged groups;

**[0062]** FIGS. 1D-G show the repeating subunit segment of exemplary morpholino oligonucleotides, designated D through G.

**[0063]** FIG. 2A shows the relative location and results of an antisense oligomer exon 51 scan designed to induce skipping of human dystrophin exon 51.

**[0064]** FIG. 2B-C shows the relative activity in cultured human rhabdomyosarcoma (RD) cells and human primary skeletal muscle cells of the three best oligomers selected from the exon 51 scan (SEQ ID NOS: 324, 326 and 327) relative to sequences (AV1-5658; SEQ ID NO: 588 and h51AON1; SEQ ID NO: 594) that are effective at inducing exon 51 skipping.

**[0065]** FIG. 2D shows the relative location within exon 51 of three selected oligomers compared to certain sequences.

**[0066]** FIG. 3A shows the relative location and results of an antisense oligomer exon 50 scan designed to induce skipping of human dystrophin exon 50 compared to other sequences that induce exon 50 skipping.

**[0067]** FIG. 3B shows the relative location and activity of antisense sequences selected from the exon 50 scan (SEQ ID NOS: 277, 287, 290 and 291) compared to other sequences (SEQ ID NOS: 584 and 585).

**[0068]** FIG. 4A shows the relative location and results of an antisense oligomer exon 53 scan designed to induce skipping of human dystrophin exon 53.

**[0069]** FIG. 4B shows the relative location of certain sequences used to compare the exon-skipping activity of those oligomers selected as being most active in the exon 53 scan.

**[0070]** FIGS. 4C-F show the results of dose-ranging studies, summarized in

**[0071]** FIG. 4G, using the oligomers selected as being most efficacious in the exon 53 scan (SEQ ID NOS: 422, 428, 429 and 431).

**[0072]** FIGS. 4H and 4I show the relative activity of certain sequences (SEQ ID NOS: 608-611) compared to the activity of the most active exon 53-skipping oligomer (SEQ ID NO: 429) in both RD cells and human primary skeletal muscle cells.

**[0073]** FIG. 5A shows the relative location and results of an antisense oligomer exon 44 scan designed to induce skipping of human dystrophin exon 44.

**[0074]** FIG. 5B shows the relative location within exon 44 of certain sequences used to compare the exon-skipping activity to those oligomers selected as being most active in the exon 44 scan.

**[0075]** FIGS. 5C-G show the results of dose-ranging studies, summarized in

**[0076]** FIG. 5H, using the oligomers selected as being most efficacious in the exon 44 scan (SEQ ID NOS: 4, 8, 11, 12 and 13).

US 2010/0130591 A1

May 27, 2010

5

[0077] FIGS. 5I and 5J show the relative activity of certain sequences (SEQ ID NOS: 600-603) compared to the activity of the most active exon 53-skipping oligomer (SEQ ID NO:12) in both RD cells and human primary skeletal muscle cells.

[0078] FIG. 6A shows the relative location and results of an antisense oligomer exon 45 scan designed to induce skipping of human dystrophin exon 45.

[0079] FIG. 6B shows the relative location within exon 45 of certain sequences used to compare the exon-skipping activity to those oligomers selected as being most active in the exon 45 scan.

[0080] FIGS. 6C-F show the results of dose-ranging studies, summarized in

[0081] FIG. 6H, using the oligomers selected as being most efficacious in the exon 45 scan (SEQ ID NOS: 27, 29, 34 and 39).

[0082] FIG. 6G uses a relatively inactive oligomer (SEQ ID NO: 49) as a negative control.

[0083] FIGS. 6I and 6J show the relative activity of certain sequences (SEQ ID NOS: 604-607) compared to the activity of the most active exon 53-skipping oligomer (SEQ ID NO: 34) in both RD cells and human primary skeletal muscle cells.

#### DETAILED DESCRIPTION OF THE INVENTION

[0084] Embodiments of the present invention relate generally to improved antisense compounds, and methods of use thereof, which are specifically designed to induce exon skipping in the dystrophin gene. Dystrophin plays a vital role in muscle function, and various muscle-related diseases are characterized by mutated forms of this gene. Hence, in certain embodiments, the improved antisense compounds described herein induce exon skipping in mutated forms of the human dystrophin gene, such as the mutated dystrophin genes found in Duchenne's muscular dystrophy (DMD) and Becker's muscular dystrophy (BMD).

[0085] Due to aberrant mRNA splicing events caused by mutations, these mutated human dystrophin genes either express defective dystrophin protein or express no measurable dystrophin at all, a condition that leads to various forms of muscular dystrophy. To remedy this condition, the antisense compounds of the present invention typically hybridize to selected regions of a pre-processed RNA of a mutated human dystrophin gene, induce exon skipping and differential splicing in that otherwise aberrantly spliced dystrophin mRNA, and thereby allow muscle cells to produce an mRNA transcript that encodes a functional dystrophin protein. In certain embodiments, the resulting dystrophin protein is not necessarily the "wild-type" form of dystrophin, but is rather a truncated, yet functional or semi-functional, form of dystrophin.

[0086] By increasing the levels of functional dystrophin protein in muscle cells, these and related embodiments may be useful in the prophylaxis and treatment of muscular dystrophy, especially those forms of muscular dystrophy, such as DMD and BMD, that are characterized by the expression of defective dystrophin proteins due to aberrant mRNA splicing. The specific oligomers described herein further provide improved, dystrophin-exon-specific targeting over other oligomers in use, and thereby offer significant and practical advantages over alternate methods of treating relevant forms of muscular dystrophy.

[0087] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly

understood by those of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, preferred methods and materials are described. For the purposes of the present invention, the following terms are defined below.

#### DEFINITIONS

[0088] The articles "a" and "an" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0089] By "about" is meant a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

[0090] By "coding sequence" is meant any nucleic acid sequence that contributes to the code for the polypeptide product of a gene. By contrast, the term "non-coding sequence" refers to any nucleic acid sequence that does not contribute to the code for the polypeptide product of a gene.

[0091] Throughout this specification, unless the context requires otherwise, the words "comprise," "comprises," and "comprising" will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

[0092] By "consisting of" is meant including, and limited to, whatever follows the phrase "consisting of." Thus, the phrase "consisting of" indicates that the listed elements are required or mandatory, and that no other elements may be present. By "consisting essentially of" is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase "consisting essentially of" indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be present depending upon whether or not they materially affect the activity or action of the listed elements.

[0093] The terms "complementary" and "complementarity" refer to polynucleotides (i.e., a sequence of nucleotides) related by the base-pairing rules. For example, the sequence "A-G-T," is complementary to the sequence "T-C-A." Complementarity may be "partial," in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or, there may be "complete" or "total" complementarity between the nucleic acids. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. While perfect complementarity is often desired, some embodiments can include one or more but preferably 6, 5, 4, 3, 2, or 1 mismatches with respect to the target RNA. Variations at any location within the oligomer are included. In certain embodiments, variations in sequence near the termini of an oligomer are generally preferable to variations in the interior, and if present are typically within about 6, 5, 4, 3, 2, or 1 nucleotides of the 5' and/or 3' terminus.

[0094] The terms "cell penetrating peptide" or "CPP" are used interchangeably and refer to cationic cell penetrating peptides, also called transport peptides, carrier peptides, or peptide transduction domains. The peptides, as shown herein,

US 2010/0130591 A1

May 27, 2010

6

have the capability of inducing cell penetration within 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of cells of a given cell culture population, including all integers in between, and allow macromolecular translocation within multiple tissues in vivo upon systemic administration.

**[0095]** The terms “antisense oligomer” or “antisense compound” are used interchangeably and refer to a sequence of cyclic subunits, each bearing a base-pairing moiety, linked by intersubunit linkages that allow the base-pairing moieties to hybridize to a target sequence in a nucleic acid (typically an RNA) by Watson-Crick base pairing, to form a nucleic acid: oligomer heteroduplex within the target sequence. The cyclic subunits are based on ribose or another pentose sugar or, in a preferred embodiment, a morpholino group (see description of morpholino oligomers below).

**[0096]** Such an antisense oligomer can be designed to block or inhibit translation of mRNA or to inhibit natural pre-mRNA splice processing, and may be said to be “directed to” or “targeted against” a target sequence with which it hybridizes. In certain embodiments, the target sequence includes a region including an AUG start codon of an mRNA, a 3' or 5' splice site of a pre-processed mRNA, or a branch point. The target sequence may be within an exon or within an intron. The target sequence for a splice site may include an mRNA sequence having its 5' end 1 to about 25 base pairs downstream of a normal splice acceptor junction in a preprocessed mRNA. A preferred target sequence for a splice is any region of a preprocessed mRNA that includes a splice site or is contained entirely within an exon coding sequence or spans a splice acceptor or donor site. An oligomer is more generally said to be “targeted against” a biologically relevant target, such as a protein, virus, or bacteria, when it is targeted against the nucleic acid of the target in the manner described above. Included are antisense oligomers that comprise, consist essentially of, or consist of one or more of SEQ ID NOS:1 to 569 and 612 to 635. Also included are variants of these antisense oligomers, including variant oligomers having 80%, 85%, 90%, 95%, 97%, 98%, or 99% (including all integers in between) sequence identity or sequence homology to any one of SEQ ID NOS:1 to 569 and 612 to 635, and/or variants that differ from these sequences by about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides, preferably those variants that induce exon skipping of one or more selected human dystrophin exons. Also included are oligomers of any one or more of SEQ ID NOS:584-611 and 634-635, which comprise a suitable number of charged linkages, as described herein, e.g. up to about 1 per every 2-5 uncharged linkages, such as about 4-5 per every 10 uncharged linkages, and/or which comprise an Arg-rich peptide attached thereto, as also described herein.

**[0097]** The terms “morpholino oligomer” or “PMO” (phosphoramidate- or phosphorodiamidate morpholino oligomer) refer to an oligonucleotide analog composed of morpholino subunit structures, where (i) the structures are linked together by phosphorus-containing linkages, one to three atoms long, preferably two atoms long, and preferably uncharged or cationic, joining the morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, and (ii) each morpholino ring bears a purine or pyrimidine base-pairing moiety effective to bind, by base specific hydrogen bonding, to a base in a polynucleotide. See, e.g., the structure in FIG. 1A, which shows a preferred phosphorodiamidate linkage type. Variations can be made to this linkage as long as they do not interfere with binding or activity. For example, the oxygen attached to phosphorus may be substituted with sulfur (thio-

phosphorodiamidate). The 5' oxygen may be substituted with amino or lower alkyl substituted amino. The pendant nitrogen attached to phosphorus may be unsubstituted, monosubstituted, or disubstituted with (optionally substituted) lower alkyl. See also the discussion of cationic linkages below. The synthesis, structures, and binding characteristics of morpholino oligomers are detailed in U.S. Pat. Nos. 5,698,685, 5,217,866, 5,142,047, 5,034,506, 5,166,315, 5,521,063, and 5,506,337, and PCT Appn. No. PCT/US07/11435 (cationic linkages), all of which are incorporated herein by reference.

**[0098]** The purine or pyrimidine base pairing moiety is typically adenine, cytosine, guanine, uracil, thymine or inosine. Also included are bases such as pyridin-4-one, pyridin-2-one, phenyl, pseudouracil, 2,4,6-trimethylthio benzene, 3-methyl uracil, dihydrouridine, naphthyl, aminophenyl, 5-alkylcytidines (e.g., 5-methylcytidine), 5-alkyluridines (e.g., ribothymidine), 5-halouridine (e.g., 5-bromouridine) or 6-azapyrimidines or 6-alkylpyrimidines (e.g. 6-methyluridine), propyne, quinosine, 2-thiouridine, 4-thiouridine, wybutosine, wybutoxosine, 4-acetyluridine, 5-(carboxyhydroxymethyl)uridine, 5"-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluridine,  $\beta$ -D-galactosylquenosine, 1-methyladenosine, 1-methylinosine, 2,2-dimethylguanosine, 3-methylcytidine, 2-methyladenosine, 2-methylguanosine, N6-methyladenosine, 7-methylguanosine, 5-methoxyaminomethyl-2-thiouridine, 5-methylaminomethyluridine, 5-methylcarbonylmethyluridine, 5-methylxyuridine, 5-methyl-2-thiouridine, 2-methylthio-N6-isopentenyladenosine,  $\beta$ -D-mannosylquenosine, uridine-5-oxyacetic acid, 2-thiocytidine, threonine derivatives and others (Burgin et al., 1996, *Biochemistry*, 35, 14090; Uhlman & Peyman, *supra*). By “modified bases” in this aspect is meant nucleotide bases other than adenine (A), guanine (G), cytosine (C), thymine (T), and uracil (U), as illustrated above; such bases can be used at any position in the antisense molecule. Persons skilled in the art will appreciate that depending on the uses of the oligomers, Ts and Us are interchangeable. For instance, with other antisense chemistries such as 2'-O-methyl antisense oligonucleotides that are more RNA-like, the T bases may be shown as U (see, e.g., Sequence ID Listing).

**[0099]** An “amino acid subunit” or “amino acid residue” can refer to an  $\alpha$ -amino acid residue (e.g.,  $-\text{CO}-\text{CHR}-\text{NH}-$ ) or a  $\beta$ - or other amino acid residue (e.g.,  $-\text{CO}-(\text{CH}_2)_n\text{CHR}-\text{NH}-$ ), where R is a side chain (which may include hydrogen) and n is 1 to 6, preferably 1 to 4.

**[0100]** The term “naturally occurring amino acid” refers to an amino acid present in proteins found in nature, such as the 20 (L)-amino acids utilized during protein biosynthesis as well as others such as 4-hydroxyproline, hydroxylysine, desmosine, isodesmosine, homocysteine, citrulline and ornithine. The term “non-natural amino acids” refers to those amino acids not present in proteins found in nature, examples include beta-alanine ( $\beta$ -Ala; or B), 6-aminohexanoic acid (Ahx) and 6-aminopentanoic acid. Additional examples of “non-natural amino acids” include, without limitation, (D)-amino acids, norleucine, norvaline, p-fluorophenylalanine, ethionine and the like, which are known to a person skilled in the art.

**[0101]** An “effective amount” or “therapeutically effective amount” refers to an amount of therapeutic compound, such as an antisense oligomer, administered to a mammalian subject, either as a single dose or as part of a series of doses, which is effective to produce a desired physiological response



US 2010/0130591 A1

May 27, 2010

7

or therapeutic effect in the subject. One example of a desired physiological response includes increased expression of a relatively functional or biologically active form of the dystrophin protein, mainly in muscle tissues or cells that contain a defective dystrophin protein or no dystrophin, as compared to no antisense oligomer or a control oligomer. Examples of desired therapeutic effects include, without limitation, improvements in the symptoms or pathology of muscular dystrophy, reducing the progression of symptoms or pathology of muscular dystrophy, and slowing the onset of symptoms or pathology of muscular dystrophy, among others. Examples of such symptoms include fatigue, mental retardation, muscle weakness, difficulty with motor skills (e.g., running, hopping, jumping), frequent falls, and difficulty walking. The pathology of muscular dystrophy can be characterized, for example, by muscle fibre damage and membrane leakage. For an antisense oligomer, this effect is typically brought about by altering the splice-processing of a selected target sequence (e.g., dystrophin), such as to induce exon skipping.

**[0102]** An “exon” refers to a defined section of nucleic acid that encodes for a protein, or a nucleic acid sequence that is represented in the mature form of an RNA molecule after either portions of a pre-processed (or precursor) RNA have been removed by splicing. The mature RNA molecule can be a messenger RNA (mRNA) or a functional form of a non-coding RNA, such as rRNA or tRNA. The human dystrophin gene has about 75 exons.

**[0103]** An “intron” refers to a nucleic acid region (within a gene) that is not translated into a protein. An intron is a non-coding section that is transcribed into a precursor mRNA (pre-mRNA), and subsequently removed by splicing during formation of the mature RNA.

**[0104]** “Exon skipping” refers generally to the process by which an entire exon, or a portion thereof, is removed from a given pre-processed RNA, and is thereby excluded from being present in the mature RNA, such as the mature mRNA that is translated into a protein. Hence, the portion of the protein that is otherwise encoded by the skipped exon is not present in the expressed form of the protein, typically creating an altered, though still functional, form of the protein. In certain embodiments, the exon being skipped is an aberrant exon from the human dystrophin gene, which may contain a mutation or other alteration in its sequence that otherwise causes aberrant splicing. In certain embodiments, the exon being skipped is any one or more of exons 1-75 of the dystrophin gene, though any one or more of exons 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, and/or 55 of the human dystrophin gene are preferred.

**[0105]** “Dystrophin” is a rod-shaped cytoplasmic protein, and a vital part of the protein complex that connects the cytoskeleton of a muscle fiber to the surrounding extracellular matrix through the cell membrane. Dystrophin contains multiple functional domains. For instance, dystrophin contains an actin binding domain at about amino acids 14-240 and a central rod domain at about amino acids 253-3040. This large central domain is formed by 24 spectrin-like triple-helical elements of about 109 amino acids, which have homology to alpha-actinin and spectrin. The repeats are typically interrupted by four proline-rich non-repeat segments, also referred to as hinge regions. Repeats 15 and 16 are separated by an 18 amino acid stretch that appears to provide a major site for proteolytic cleavage of dystrophin. The sequence identity between most repeats ranges from 10-25%. One repeat con-

tains three alpha-helices: 1, 2 and 3. Alpha-helices 1 and 3 are each formed by 7 helix turns, probably interacting as a coiled-coil through a hydrophobic interface. Alpha-helix 2 has a more complex structure and is formed by segments of four and three helix turns, separated by a Glycine or Proline residue. Each repeat is encoded by two exons, typically interrupted by an intron between amino acids 47 and 48 in the first part of alpha-helix 2. The other intron is found at different positions in the repeat, usually scattered over helix-3. Dystrophin also contains a cysteine-rich domain at about amino acids 3080-3360), including a cysteine-rich segment (i.e., 15 Cysteines in 280 amino acids) showing homology to the C-terminal domain of the slime mold (*Dictyostelium discoideum*) alpha-actinin. The carboxy-terminal domain is at about amino acids 3361-3685.

**[0106]** The amino-terminus of dystrophin binds to F-actin and the carboxy-terminus binds to the dystrophin-associated protein complex (DAPC) at the sarcolemma. The DAPC includes the dystroglycans, sarcoglycans, integrins and caveolin, and mutations in any of these components cause autosomally inherited muscular dystrophies. The DAPC is destabilized when dystrophin is absent, which results in diminished levels of the member proteins, and in turn leads to progressive fibre damage and membrane leakage. In various forms of muscular dystrophy, such as Duchenne’s muscular dystrophy (DMD) and Becker’s muscular dystrophy (BMD), muscle cells produce an altered and functionally defective form of dystrophin, or no dystrophin at all, mainly due to mutations in the gene sequence that lead to incorrect splicing. The predominant expression of the defective dystrophin protein, or the complete lack of dystrophin or a dystrophin-like protein, leads to rapid progression of muscle degeneration, as noted above. In this regard, a “defective” dystrophin protein may be characterized by the forms of dystrophin that are produced in certain subjects with DMD or BMD, as known in the art, or by the absence of detectable dystrophin.

**[0107]** Table A provides an illustration of the various dystrophin domains, the amino acid residues that encompass these domains, and the exons that encode them.

TABLE A

Domain	Sub Domain	Residue Nos	Exons
actin binding domain		14-240	2-8
central rod domain		253-3040	8-61
	hinge 1	253-327	(8)-9
	repeat 1	337-447	10-11
	repeat 2	448-556	12-14
	repeat 3	557-667	14-16
	hinge 2	668-717	17
	repeat 4	718-828	(17)-20
	repeat 5	829-934	20-21
	repeat 6	935-1045	22-23
	repeat 7	1046-1154	(23)-(26)
	repeat 8	1155-1263	26-27
	repeat 9	1264-1367	28-(30)
	repeat 10	1368-1463	30-32
	repeat 11	1464-1568	32-(34)
	repeat 12	1569-1676	34-35
	repeat 13	1677-1778	36-37
	repeat 14	1779-1874	38-(40)
	repeat 15	1875-1973	40-41
	interruption	1974-1991	42
	repeat 16	1992-2101	42-43
	repeat 17	2102-2208	44-45



US 2010/0130591 A1

May 27, 2010

8

TABLE A-continued

Domain	Sub Domain	Residue Nos	Exons
Cysteine-rich domain	repeat 18	2209-2318	46-48
	repeat 19	2319-2423	48-50
	hinge 3	2424-2470	50-51
	repeat 20	2471-2577	51-53
	repeat 21	2578-2686	53-(55)
	repeat 22	2687-2802	55-(57)
	repeat 23	2803-2931	57-59
	repeat 24	2932-3040	59-(61)
	hinge 4	3041-3112	61-64
		3080-3360	63-69
Carboxy-terminal domain	dystroglycan binding site	3080-3408	63-70
	WW domain	3056-3092	62-63
	EF-hand 1	3130-3157	65
	EF-hand 2	3178-3206	65-66
	ZZ domain	3307-3354	68-69
		3361-3685	70-79
	alpha1-syntrophin binding site	3444-3494	73-74
	β1-syntrophin binding site (Leu)6-heptad repeat	3495-3535 3558-3593	74-75 75

**[0108]** As used herein, the terms “function” and “functional” and the like refer to a biological, enzymatic, or therapeutic function.

**[0109]** A “functional” dystrophin protein refers generally to a dystrophin protein having sufficient biological activity to reduce the progressive degradation of muscle tissue that is otherwise characteristic of muscular dystrophy, typically as compared to the altered or “defective” form of dystrophin protein that is present in certain subjects with DMD or BMD. In certain embodiments, a functional dystrophin protein may have about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% (including all integers in between) of the in vitro or in vivo biological activity of wild-type dystrophin, as measured according to routine techniques in the art. As one example, dystrophin-related activity in muscle cultures in vitro can be measured according to myotube size, myofibril organization (or disorganization), contractile activity, and spontaneous clustering of acetylcholine receptors (see, e.g., Brown et al., *Journal of Cell Science*, 112:209-216, 1999). Animal models are also valuable resources for studying the pathogenesis of disease, and provide a means to test dystrophin-related activity. Two of the most widely used animal models for DMD research are the mdx mouse and the golden retriever muscular dystrophy (GRMD) dog, both of which are dystrophin negative (see, e.g., Collins & Morgan, *Int J Exp Pathol* 84: 165-172, 2003). These and other animal models can be used to measure the functional activity of various dystrophin proteins. Included are truncated forms of dystrophin, such as those forms that are produced by certain of the exon-skipping antisense compounds of the present invention.

**[0110]** By “gene” is meant a unit of inheritance that occupies a specific locus on a chromosome and consists of transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (i.e., introns, 5' and 3' untranslated sequences).

**[0111]** By “isolated” is meant material that is substantially or essentially free from components that normally accompany it in its native state. For example, an “isolated polynucleotide,” as used herein, may refer to a polynucleotide that has been purified or removed from the sequences that flank it in a

naturally-occurring state, e.g., a DNA fragment that has been removed from the sequences that are normally adjacent to the fragment.

**[0112]** By “enhance” or “enhancing,” or “increase” or “increasing,” or “stimulate” or “stimulating,” refers generally to the ability of one or antisense compounds or compositions to produce or cause a greater physiological response (i.e., downstream effects) in a cell or a subject, as compared to the response caused by either no antisense compound or a control compound. A measurable physiological response may include increased expression of a functional form of a dystrophin protein, or increased dystrophin-related biological activity in muscle tissue, among other responses apparent from the understanding in the art and the description herein. Increased muscle function can also be measured, including increases or improvements in muscle function by about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%. The percentage of muscle fibres that express a functional dystrophin can also be measured, including increased dystrophin expression in about 1%, 2%, %, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of muscle fibres. For instance, it has been shown that around 40% of muscle function improvement can occur if 25-30% of fibers express dystrophin (see, e.g., DeloRusso et al, *Proc Natl Acad Sci USA* 99: 12979-12984, 2002). An “increased” or “enhanced” amount is typically a “statistically significant” amount, and may include an increase that is 1.1, 1.2, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1), e.g., 1.5, 1.6, 1.7, 1.8, etc.) the amount produced by no antisense compound (the absence of an agent) or a control compound.

**[0113]** The term “reduce” or “inhibit” may relate generally to the ability of one or more antisense compounds of the invention to “decrease” a relevant physiological or cellular response, such as a symptom of a disease or condition described herein, as measured according to routine techniques in the diagnostic art. Relevant physiological or cellular responses (in vivo or in vitro) will be apparent to persons skilled in the art, and may include reductions in the symptoms or pathology of muscular dystrophy, or reductions in the expression of defective forms of dystrophin, such as the altered forms of dystrophin that are expressed in individuals with DMD or BMD. A “decrease” in a response may be statistically significant as compared to the response produced by no antisense compound or a control composition, and may include a 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% decrease, including all integers in between.

**[0114]** “Homology” refers to the percentage number of amino acids that are identical or constitute conservative substitutions. Homology may be determined using sequence comparison programs such as GAP (Deveraux et al., 1984, *Nucleic Acids Research* 12, 387-395). In this way sequences of a similar or substantially different length to those cited herein could be compared by insertion of gaps into the alignment, such gaps being determined, for example, by the comparison algorithm used by GAP.

US 2010/0130591 A1

May 27, 2010

9

**[0115]** The recitations “sequence identity” or, for example, comprising a “sequence 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

**[0116]** Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include “reference sequence,” “comparison window,” “sequence identity,” “percentage of sequence identity,” and “substantial identity.” A “reference sequence” is at least 8 or 10 but frequently 15 to 18 and often at least 25 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e., only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a “comparison window” to identify and compare local regions of sequence similarity. A “comparison window” refers to a conceptual segment of at least 6 contiguous positions, usually about 50 to about 100, more usually about 100 to about 150 in which a sequence is compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. The comparison window may comprise additions or deletions (i.e., gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, Wis., USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., 1997, *Nucl. Acids Res.* 25:3389. A detailed discussion of sequence analysis can be found in Unit 19.3 of Ausubel et al., “Current Protocols in Molecular Biology,” John Wiley & Sons Inc, 1994-1998, Chapter 15.

**[0117]** “Treatment” or “treating” of an individual (e.g., a mammal, such as a human) or a cell may include any type of intervention used in an attempt to alter the natural course of the individual or cell. Treatment includes, but is not limited to, administration of a pharmaceutical composition, and may be performed either prophylactically or subsequent to the initiation of a pathologic event or contact with an etiologic agent. Treatment includes any desirable effect on the symptoms or pathology of a disease or condition associated with the dystrophin protein, as in certain forms of muscular dystrophy,

and may include, for example, minimal changes or improvements in one or more measurable markers of the disease or condition being treated. Also included are “prophylactic” treatments, which can be directed to reducing the rate of progression of the disease or condition being treated, delaying the onset of that disease or condition, or reducing the severity of its onset. “Treatment” or “prophylaxis” does not necessarily indicate complete eradication, cure, or prevention of the disease or condition, or associated symptoms thereof.

**[0118]** Hence, included are methods of treating muscular dystrophy, such as DMD and BMD, by administering one or more antisense oligomers of the present invention (e.g., SEQ ID NOS: 1 to 569 and 612 to 635, and variants thereof), optionally as part of a pharmaceutical formulation or dosage form, to a subject in need thereof. Also included are methods of inducing exon-skipping in a subject by administering one or more antisense oligomers, in which the exon is one of exons 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, and/or 55 from the dystrophin gene, preferably the human dystrophin gene. A “subject,” as used herein, includes any animal that exhibits a symptom, or is at risk for exhibiting a symptom, which can be treated with an antisense compound of the invention, such as a subject that has or is at risk for having DMD or BMD, or any of the symptoms associated with these conditions (e.g., muscle fibre loss). Suitable subjects (patients) include laboratory animals (such as mouse, rat, rabbit, or guinea pig), farm animals, and domestic animals or pets (such as a cat or dog). Non-human primates and, preferably, human patients, are included.

**[0119]** Also included are vector delivery systems that are capable of expressing the oligomeric, dystrophin-targeting sequences of the present invention, such as vectors that express a polynucleotide sequence comprising any one or more of SEQ ID NOS: 1 to 569 and 612 to 635, or variants thereof, as described herein. By “vector” or “nucleic acid construct” is meant a polynucleotide molecule, preferably a DNA molecule derived, for example, from a plasmid, bacteriophage, yeast or virus, into which a polynucleotide can be inserted or cloned. A vector preferably contains one or more unique restriction sites and can be capable of autonomous replication in a defined host cell including a target cell or tissue or a progenitor cell or tissue thereof, or be integrable with the genome of the defined host such that the cloned sequence is reproducible. Accordingly, the vector can be an autonomously replicating vector, i.e., a vector that exists as an extra-chromosomal entity, the replication of which is independent of chromosomal replication, e.g., a linear or closed circular plasmid, an extra-chromosomal element, a mini-chromosome, or an artificial chromosome. The vector can contain any means for assuring self-replication. Alternatively, the vector can be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated.

**[0120]** A vector or nucleic acid construct system can comprise a single vector or plasmid, two or more vectors or plasmids, which together contain the total DNA to be introduced into the genome of the host cell, or a transposon. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. In the present case, the vector or nucleic acid construct is preferably one which is operably functional in a mammalian cell, such as a muscle cell. The vector can also include a selection marker such as an antibiotic or drug resistance gene, or a reporter gene (i.e., green fluorescent protein,

US 2010/0130591 A1

May 27, 2010

10

luciferase), that can be used for selection or identification of suitable transformants or transfectants. Exemplary delivery systems may include viral vector systems (i.e., viral-mediated transduction) including, but not limited to, retroviral (e.g., lentiviral) vectors, adenoviral vectors, adeno-associated viral vectors, and herpes viral vectors, among others known in the art.

**[0121]** The term “operably linked” as used herein means placing an oligomer-encoding sequence under the regulatory control of a promoter, which then controls the transcription of the oligomer.

**[0122]** A wild-type gene or gene product is that which is most frequently observed in a population and is thus arbitrarily designed the “normal” or “wild-type” form of the gene.

**[0123]** “Alkyl” or “alkylene” both refer to a saturated straight or branched chain hydrocarbon radical containing from 1 to 18 carbons. Examples include without limitation methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl and n-hexyl. The term “lower alkyl” refers to an alkyl group, as defined herein, containing between 1 and 8 carbons.

**[0124]** “Alkenyl” refers to an unsaturated straight or branched chain hydrocarbon radical containing from 2 to 18 carbons and comprising at least one carbon to carbon double bond. Examples include without limitation ethenyl, propenyl, iso-propenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl and n-hexenyl. The term “lower alkenyl” refers to an alkenyl group, as defined herein, containing between 2 and 8 carbons.

**[0125]** “Alkynyl” refers to an unsaturated straight or branched chain hydrocarbon radical containing from 2 to 18 carbons comprising at least one carbon to carbon triple bond. Examples include without limitation ethynyl, propynyl, isopropynyl, butynyl, iso-butynyl, tert-butynyl, pentynyl and hexynyl. The term “lower alkynyl” refers to an alkynyl group, as defined herein, containing between 2 and 8 carbons.

**[0126]** “Cycloalkyl” refers to a mono- or poly-cyclic alkyl radical. Examples include without limitation cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

**[0127]** “Aryl” refers to a cyclic aromatic hydrocarbon moiety containing from 6 to 18 carbons having one or more closed ring(s). Examples include without limitation phenyl, benzyl, naphthyl, anthracenyl, phenanthracenyl and biphenyl.

**[0128]** “Aralkyl” refers to a radical of the formula RaRb where Ra is an alkylene chain as defined above and Rb is one or more aryl radicals as defined above, for example, benzyl, diphenylmethyl and the like.

**[0129]** “Thioalkoxy” refers to a radical of the formula —SRc where Rc is an alkyl radical as defined herein. The term “lower thioalkoxy” refers to an alkoxy group, as defined herein, containing between 1 and 8 carbons.

**[0130]** “Alkoxy” refers to a radical of the formula —ORd where Rd is an alkyl radical as defined herein. The term “lower alkoxy” refers to an alkoxy group, as defined herein, containing between 1 and 8 carbons. Examples of alkoxy groups include, without limitation, methoxy and ethoxy.

**[0131]** “Alkoxyalkyl” refers to an alkyl group substituted with an alkoxy group.

**[0132]** “Carbonyl” refers to the C(=O)— radical.

**[0133]** “Guanidynyl” refers to the H<sub>2</sub>N(C=NH<sub>2</sub>)—NH— radical.

**[0134]** “Amidynyl” refers to the H<sub>2</sub>N(C=NH<sub>2</sub>)CH— radical.

**[0135]** “Amino” refers to the NH<sub>2</sub> radical.

**[0136]** “Alkylamino” refers to a radical of the formula —NHRd or —NRdRd where each Rd is, independently, an alkyl radical as defined herein. The term “lower alkylamino” refers to an alkylamino group, as defined herein, containing between 1 and 8 carbons.

**[0137]** “Heterocycle” means a 5- to 7-membered monocyclic, or 7- to 10-membered bicyclic, heterocyclic ring which is either saturated, unsaturated, or aromatic, and which contains from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen heteroatom may be optionally quaternized, including bicyclic rings in which any of the above heterocycles are fused to a benzene ring. The heterocycle may be attached via any heteroatom or carbon atom. Heterocycles include heteroaryls as defined below. Thus, in addition to the heteroaryls listed below, heterocycles also include morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperiziny, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydropyrimidinyl, tetrahydrothiopyranyl, and the like.

**[0138]** “Heteroaryl” means an aromatic heterocycle ring of 5- to 10 members and having at least one heteroatom selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom, including both mono- and bicyclic ring systems. Representative heteroaryls are pyridyl, furyl, benzofuranyl, thiophenyl, benzothiophenyl, quinolinyl, pyrrolyl, indolyl, oxazolyl, benzoxazolyl, imidazolyl, benzimidazolyl, thiazolyl, benzothiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, cinnolinyl, phthalazinyl, and quinazolinyl.

**[0139]** The terms “optionally substituted alkyl”, “optionally substituted alkenyl”, “optionally substituted alkoxy”, “optionally substituted thioalkoxy”, “optionally substituted alkyl amino”, “optionally substituted lower alkyl”, “optionally substituted lower alkenyl”, “optionally substituted lower alkoxy”, “optionally substituted lower thioalkoxy”, “optionally substituted lower alkyl amino” and “optionally substituted heterocyclyl” mean that, when substituted, at least one hydrogen atom is replaced with a substituent. In the case of an oxo substituent (=O) two hydrogen atoms are replaced. In this regard, substituents include: deuterium, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heterocycle, optionally substituted cycloalkyl, oxo, halogen, —CN, —ORx, NRxRy, NRxC(=O)Ry, NRxSO<sub>2</sub>Ry, —NRxC(=O)NRxRy, C(=O)Rx, C(=O)ORx, C(=O)NRxRy, —SOMRx and —SOMNRxRy, wherein m is 0, 1 or 2, Rx and Ry are the same or different and independently hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heterocycle or optionally substituted cycloalkyl and each of said optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heterocycle and optionally substituted cycloalkyl substituents may be further substituted with one or more of oxo, halogen, —CN, —ORx, NRxRy, NRxC(=O)Ry, NRxSO<sub>2</sub>Ry, —NRxC(=O)NRxRy, C(=O)Rx, C(=O)ORx, C(=O)NRxRy, —SOMRx and —SOMNRxRy.

Constructing Antisense Oligonucleotides

**[0140]** Examples of morpholino oligonucleotides having phosphorus-containing backbone linkages are illustrated in



US 2010/0130591 A1

May 27, 2010

11

FIGS. 1A-1C. Especially preferred is a phosphorodiamidate-linked morpholino oligonucleotide such as shown in FIG. 1C, which is modified, in accordance with one aspect of the present invention, to contain positively charged groups at preferably 10%-50% of its backbone linkages. Morpholino oligonucleotides with uncharged backbone linkages and their preparation, including antisense oligonucleotides, are detailed, for example, in (Summerton and Weller 1997) and in co-owned U.S. Pat. Nos. 5,698,685, 5,217,866, 5,142,047, 5,034,506, 5,166,315, 5,185,444, 5,521,063, and 5,506,337, all of which are expressly incorporated by reference herein.

[0141] Important properties of the morpholino-based subunits include: 1) the ability to be linked in an oligomeric form by stable, uncharged or positively charged backbone linkages; 2) the ability to support a nucleotide base (e.g. adenine, cytosine, guanine, thymidine, uracil and inosine) such that the polymer formed can hybridize with a complementary-base target nucleic acid, including target RNA,  $T_m$  values above about 45° C. in relatively short oligonucleotides (e.g., 10-15 bases); 3) the ability of the oligonucleotide to be actively or passively transported into mammalian cells; and 4) the ability of the antisense oligonucleotide:RNA heteroduplex to resist RNase and RNaseH degradation, respectively.

[0142] Exemplary backbone structures for antisense oligonucleotides of the claimed subject matter include the morpholino subunit types shown in FIGS. 1D-G, each linked by an uncharged or positively charged, phosphorus-containing subunit linkage. FIG. 1D shows a phosphorus-containing linkage which forms the five atom repeating-unit backbone, wherein the morpholino rings are linked by a 1-atom phosphoamide linkage. FIG. 1E shows a linkage which produces a 6-atom repeating-unit backbone. In this structure, the atom Y linking the 5' morpholino carbon to the phosphorus group may be sulfur, nitrogen, carbon or, preferably, oxygen. The X moiety pendant from the phosphorus may be fluorine, an alkyl or substituted alkyl, an alkoxy or substituted alkoxy, a thioalkoxy or substituted thioalkoxy, or unsubstituted, mono-substituted, or disubstituted nitrogen, including cyclic structures, such as morpholines or piperidines. Alkyl, alkoxy and thioalkoxy preferably include 1-6 carbon atoms. The Z moieties are sulfur or oxygen, and are preferably oxygen.

[0143] The linkages shown in FIGS. 1F and 1G are designed for 7-atom unit-length backbones. In structure 1F, the X moiety is as in Structure 1E, and the Y moiety may be methylene, sulfur, or, preferably, oxygen. In Structure 1G, the X and Y moieties are as in Structure 1E. Particularly preferred morpholino oligonucleotides include those composed of morpholino subunit structures of the form shown in FIG. 1E, where  $X=NH_2$ ,  $N(CH_3)_2$ , optionally substituted 1-piperazinyl, or other charged group,  $Y=O$ , and  $Z=O$ .

[0144] As noted above, the uncharged or substantially uncharged oligonucleotide may be modified, in accordance with an aspect of the invention, to include charged linkages, e.g. up to about 1 per every 2-5 uncharged linkages, such as about 4-5 per every 10 uncharged linkages. Optimal improvement in antisense activity may be seen when about 25% of the backbone linkages are cationic, including about 20% to about 30%. Also included are oligomers in which about 35%, 40%, 45%, 50%, 55%, 60% (including all integers in between), or more of the backbone linkages are cationic. Enhancement is also seen with a small number, e.g., 5% or 10-20%, of cationic linkages.


[0145] A substantially uncharged, phosphorus containing backbone in an oligonucleotide analog is typically one in

which a majority of the subunit linkages, e.g., between 50%-100%, typically at least 60% to 100% or 75% or 80% of its linkages, are uncharged at physiological pH and contain a single phosphorous atom.

[0146] Additional experiments conducted in support of the present invention indicate that the enhancement seen with added cationic backbone charges may, in some cases, be further enhanced by distributing the bulk of the charges close to the "center-region" backbone linkages of the antisense oligonucleotide, e.g., in a 20mer oligonucleotide with 8 cationic backbone linkages, having at least 70% of these charged linkages localized in the 10 centermost linkages.

[0147] The antisense compounds can be prepared by step-wise solid-phase synthesis, employing methods detailed in the references cited above, and below with respect to the synthesis of oligonucleotides having a mixture of uncharged and cationic backbone linkages. In some cases, it may be desirable to add additional chemical moieties to the antisense compound, e.g. to enhance pharmacokinetics or to facilitate capture or detection of the compound. Such a moiety may be covalently attached, typically to a terminus of the oligomer, according to standard synthetic methods. For example, addition of a polyethyleneglycol moiety or other hydrophilic polymer, e.g., one having 10-100 monomeric subunits, may be useful in enhancing solubility. One or more charged groups, e.g., anionic charged groups such as an organic acid, may enhance cell uptake. A reporter moiety, such as fluorescein or a radiolabeled group, may be attached for purposes of detection. Alternatively, the reporter label attached to the oligomer may be a ligand, such as an antigen or biotin, capable of binding a labeled antibody or streptavidin. In selecting a moiety for attachment or modification of an antisense compound, it is generally of course desirable to select chemical compounds of groups that are biocompatible and likely to be tolerated by a subject without undesirable side effects.

[0148] As noted above, the antisense compound can be constructed to contain a selected number of cationic linkages interspersed with uncharged linkages of the type described above. The intersubunit linkages, both uncharged and cationic, preferably are phosphorus-containing linkages, having the structure (II):



wherein:

[0149] W is —S— or —O—, and is preferably —O—,

[0150]  $X=NR^1R^2$  or  $OR^6$ ,

[0151]  $Y=O$  or  $NR^7$ , and

[0152] each said linkage in the oligomer is selected from:

[0153] (a) an uncharged linkage (a), wherein each of  $R^1$ ,  $R^2$ ,  $R^6$  and  $R^7$  is independently selected from hydrogen and lower alkyl;


**[0166]** Preferably, all cationic linkages in the oligomer are of the same type; i.e. all of type (b1), all of type (b2), or all of type (b3).

US 2010/0130591 A1

May 27, 2010

13

[0167] In further embodiments, the cationic linkages are selected from linkages (b1') and (b1'') as shown below, where (b1') is referred to herein as a "Pip" linkage and (b1'') is referred to herein as a "GuX" linkage:



[0168] In the structures above, W is S or O, and is preferably O; each of R<sup>1</sup> and R<sup>2</sup> is independently selected from hydrogen and optionally substituted lower alkyl, and is preferably methyl; and A represents hydrogen or a non-interfering substituent (i.e. a substituent that does not adversely affect the ability of an oligomer to bind to its intended target) on one or more carbon atoms in (b1') and (b1''). Preferably, the ring carbons in the piperazine ring are unsubstituted; however, the ring carbons of the piperazine ring may include non-interfering substituents, such as methyl or fluorine. Preferably, at most one or two carbon atoms is so substituted.

[0169] In further embodiments, at least 10% of the linkages are of type (b1') or (b1''); for example, 10%-60% and preferably 20% to 50%, of the linkages may be of type (b1') or (b1'').

[0170] In other embodiments, the oligomer contains no linkages of the type (b1') above. Alternatively, the oligomer contains no linkages of type (b1) where each R is H, R<sup>3</sup> is H or —CH<sub>3</sub>, and R<sup>4</sup> is H, —CH<sub>3</sub>, or an electron pair.

[0171] The morpholino subunits may also be linked by non-phosphorus-based intersubunit linkages, as described further below, where at least one linkage is modified with a pendant cationic group as described above.

[0172] Other oligonucleotide analog linkages which are uncharged in their unmodified state but which could also bear a pendant amine substituent could be used. For example, a 5' nitrogen atom on a morpholino ring could be employed in a sulfamide linkage or a urea linkage (where phosphorus is replaced with carbon or sulfur, respectively) and modified in a manner analogous to the 5'-nitrogen atom in structure (b3) above.

[0173] Oligomers having any number of cationic linkages are provided, including fully cationic-linked oligomers. Preferably, however, the oligomers are partially charged, having, for example, 10%-80%. In preferred embodiments, about 10% to 60%, and preferably 20% to 50% of the linkages are cationic.

[0174] In one embodiment, the cationic linkages are interspersed along the backbone. The partially charged oligomers preferably contain at least two consecutive uncharged link-

ages; that is, the oligomer preferably does not have a strictly alternating pattern along its entire length.

[0175] Also considered are oligomers having blocks of cationic linkages and blocks of uncharged linkages; for example, a central block of uncharged linkages may be flanked by blocks of cationic linkages, or vice versa. In one embodiment, the oligomer has approximately equal-length 5', 3' and center regions, and the percentage of cationic linkages in the center region is greater than about 50%, preferably greater than about 70%.

[0176] Oligomers for use in antisense applications generally range in length from about 10 to about 40 subunits, more preferably about 10 to 30 subunits, and typically 15-25 bases, including those having 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 bases. In certain embodiments, an oligomer of the invention having 19-20 subunits, a useful length for an antisense compound, may ideally have two to ten, e.g., four to eight, cationic linkages, and the remainder uncharged linkages. An oligomer having 14-15 subunits may ideally have two to seven, e.g., 3 to 5, cationic linkages, and the remainder uncharged linkages.

[0177] Each morpholino ring structure supports a base pairing moiety, to form a sequence of base pairing moieties which is typically designed to hybridize to a selected antisense target in a cell or in a subject being treated. The base pairing moiety may be a purine or pyrimidine found in native DNA or RNA (e.g., A, G, C, T or U) or an analog, such as hypoxanthine (the base component of the nucleoside inosine) or 5-methyl cytosine.

#### Peptide Transporters

[0178] The antisense compounds of the invention may include an oligonucleotide moiety conjugated to an arginine-rich peptide transport moiety effective to enhance transport of the compound into cells. The transport moiety is preferably attached to a terminus of the oligomer, as shown, for example, in FIGS. 1B and 1C. The peptide transport moiety preferably comprises 6 to 16 subunits selected from X' subunits, Y' subunits, and Z' subunits, wherein:

[0179] (a) each X' subunit independently represents lysine, arginine or an arginine analog, said analog being a cationic α-amino acid comprising a side chain of the structure R<sup>1</sup>N=C(NH<sub>2</sub>)R<sup>2</sup>, where R<sup>1</sup> is H or R; R<sup>2</sup> is R, —NH<sub>2</sub>, —NHR, or —NR<sub>2</sub>, where R is optionally substituted lower alkyl or optionally substituted lower alkenyl; R<sup>1</sup> and R<sup>2</sup> may join together to form a ring; and the side chain is linked to said amino acid via R<sup>1</sup> or R<sup>2</sup>;

[0180] (b) each Y' subunit independently represents a neutral amino acid —C(=O)—(CHR)<sub>n</sub>—NH—, where n is 2 to 7 and each R is independently H or methyl; and

[0181] (c) each Z' subunit independently represents an α-amino acid having a neutral aralkyl side chain; wherein the peptide comprises a sequence represented by at least one of (X'Y'X')<sub>p</sub>, (X'Y')<sub>m</sub>, and/or (X'Z'Z')<sub>p</sub>, where p is 2 to 5 and m is 2 to 8. Certain embodiments include various combinations selected independently from (X'Y'X')<sub>p</sub>, (X'Y')<sub>m</sub>, and/or (X'Z'Z')<sub>p</sub>, including, for example, peptides having the sequence (X'Y'X')(X'Z'Z')(X'Y'X')(X'Z'Z') (SEQ ID NO:637).

[0182] In selected embodiments, for each X', the side chain moiety is guanidyl, as in the amino acid subunit arginine (Arg). In certain embodiments, each Y' is independently —C(=O)—(CH<sub>2</sub>)<sub>n</sub>—CHR—NH—, where n is 2 to 7 and R is

US 2010/0130591 A1

May 27, 2010

14

H. For example, when n is 5 and R is H, Y' is a 6-amino-hexanoic acid subunit, abbreviated herein as Ahx; when n is 2 and R is H, Y' is a  $\beta$ -alanine subunit, abbreviated herein as B. Certain embodiments relate to carrier peptides having a combination of different neutral amino acids, including, for example, peptides comprising the sequence —RahxRRBRRAhxRRBRRAhxB— (SEQ ID NO:578), which contains both  $\beta$ -alanine and 6-aminohexanoic acid.

[0183] Preferred peptides of this type include those comprising arginine dimers alternating with single Y' subunits, where Y' is preferably Ahx or B or both. Examples include peptides having the formula  $(RY'R)_p$  and/or the formula  $(RRY')_p$ , where p is 1 to 2 to 5 and where Y' is preferably Ahx. In one embodiment, Y' is a 6-aminohexanoic acid subunit, R is arginine and p is 4. Certain embodiments include various linear combinations of at least two of  $(RY'R)_p$  and  $(RRY')_p$ , including, for example, illustrative peptides having the sequence  $(RY'R)(RRY')(RY'R)(RRY')$  (SEQ ID NO:638), or  $(RRY')(RY'R)(RRY')$  (SEQ ID NO:639). Other combinations are contemplated. In a further illustrative embodiment, each Z' is phenylalanine, and m is 3 or 4.

[0184] The conjugated peptide is preferably linked to a terminus of the oligomer via a linker Ahx-B, where Ahx is a 6-aminohexanoic acid subunit and B is a  $\beta$ -alanine subunit, as shown, for example, in FIGS. 1B and 1C.

[0185] In selected embodiments, for each X', the side chain moiety is independently selected from the group consisting of guanidyl ( $\text{HN}=\text{C}(\text{NH}_2)\text{NH}-$ ), amidinyl ( $\text{HN}=\text{C}(\text{NH}_2)\text{CH}-$ ), 2-aminodihydropyrimidyl, 2-aminotetrahydropyrimidyl, 2-aminopyridinyl, and 2-aminopyrimidinyl, and it is preferably selected from guanidyl and amidinyl. In one embodiment, the side chain moiety is guanidyl, as in the amino acid subunit arginine (Arg).

[0186] In certain embodiments, the Y' subunits may be contiguous, in that no X' subunits intervene between Y' subunits, or interspersed singly between X' subunits. In certain embodiments, the linking subunit may be between Y' subunits. In one embodiment, the Y' subunits are at a terminus of the transporter; in other embodiments, they are flanked by X' subunits. In further preferred embodiments, each Y' is  $-\text{C}(=\text{O})-(\text{CH}_2)_n-\text{CHR}-\text{NH}-$ , where n is 2 to 7 and R is H. For example, when n is 5 and R is H, Y' is a 6-amino-hexanoic acid subunit, abbreviated herein as Ahx. In selected embodiments of this group, each X' comprises a guanidyl side chain moiety, as in an arginine subunit. Preferred peptides of this type include those comprising arginine dimers alternat-

ing with single Y' subunits, where Y' is preferably Ahx. Examples include peptides having the formula  $(RY'R)_4$  or the formula  $(RRY')_4$ , where Y' is preferably Ahx. In the latter case, the nucleic acid analog is preferably linked to a terminal Y' subunit, preferably at the C-terminus, as shown, for example, in FIGS. 1B and 1C. The preferred linker is of the structure AhxB, where Ahx is a 6-aminohexanoic acid subunit and B is a  $\beta$ -alanine subunit.

[0187] The transport moieties as described above have been shown to greatly enhance cell entry of attached oligomers, relative to uptake of the oligomer in the absence of the attached transport moiety, and relative to uptake by an attached transport moiety lacking the hydrophobic subunits Y'. Such enhanced uptake is preferably evidenced by at least a two-fold increase, and preferably a four-fold increase, in the uptake of the compound into mammalian cells relative to uptake of the agent by an attached transport moiety lacking the hydrophobic subunits Y'. Uptake is preferably enhanced at least twenty fold, and more preferably forty fold, relative to the unconjugated compound.

[0188] A further benefit of the transport moiety is its expected ability to stabilize a duplex between an antisense compound and its target nucleic acid sequence, presumably by virtue of electrostatic interaction between the positively charged transport moiety and the negatively charged nucleic acid. The number of charged subunits in the transporter is less than 14, as noted above, and preferably between 8 and 11, since too high a number of charged subunits may lead to a reduction in sequence specificity.

[0189] The use of arginine-rich peptide transporters (i.e., cell-penetrating peptides) is particularly useful in practicing the present invention. Certain peptide transporters have been shown to be highly effective at delivery of antisense compounds into primary cells including muscle cells (Marshall, Oda et al. 2007; Jearawiriyapaisarn, Moulton et al. 2008; Wu, Moulton et al. 2008). Furthermore, compared to other peptide transporters such as Penetratin and the Tat peptide, the peptide transporters described herein, when conjugated to an antisense PMO, demonstrate an enhanced ability to alter splicing of several gene transcripts (Marshall, Oda et al. 2007). Especially preferred are the P007, CP06062 and CP04057 transport peptides listed below in Table 3 (SEQ ID NOS: 573, 578 and 577, respectively).

[0190] Exemplary peptide transporters, including linkers (B or AhxB) are given below in Table B below. Preferred sequences are those designated CP06062 (SEQ ID NO: 578), P007 (SEQ ID NO: 573) and CP04057 (SEQ ID NO: 577).

TABLE B

Exemplary Peptide Transporters for Intracellular Delivery of PMO		
Peptide	Sequence (N-terminal to C-terminal)	SEQ ID NO:
rTAT	RRRQRKKRC	570
R <sub>9</sub> F <sub>2</sub>	RRRRRRRRFFC	571
(RAhx) <sub>4</sub> B	RAhxRAhxRAhxRAhxRAhxB	572
(RAhxR) <sub>4</sub> AhxB; (P007)	RAhxRAhxRAhxRAhxRAhxRAhxB	573
(AhxRR) <sub>4</sub> AhxB	AhxRAhxRAhxRAhxRAhxRAhxB	574
(RAhx) <sub>6</sub> B	RAhxRAhxRAhxRAhxRAhxRAhxB	575



TABLE B-continued

Exemplary Peptide Transporters for Intracellular Delivery of PMO		
Peptide	Sequence (N-terminal to C-terminal)	SEQ ID NO:
(RAhx) <sub>8</sub> B	RAhxRAhxRAhxRAhxRAhxRAhxRAhxB	576
(RAhxR) <sub>5</sub> AhxB (CP05057)	RAhxRRAhxRRAhxRRAhxRRAhxRAhxB	577
(RAhxRRBR) <sub>2</sub> AhxB; (CP06062)	RAhxRRBRRAhxRRBRAhxB	578
MSP	ASSLNIA	579

## Formulations

**[0191]** In certain embodiments, the present invention provides formulations or compositions suitable for the therapeutic delivery of antisense oligomers, as described herein. Hence, in certain embodiments, the present invention provides pharmaceutically acceptable compositions that comprise a therapeutically-effective amount of one or more of the oligomers described herein, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. While it is possible for an oligomer of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation (composition).

**[0192]** Methods for the delivery of nucleic acid molecules are described, for example, in Akhtar et al., 1992, *Trends Cell Bio.*, 2:139; and *Delivery Strategies for Antisense Oligonucleotide Therapeutics*, ed. Akhtar; Sullivan et al., PCT WO 94/02595. These and other protocols can be utilized for the delivery of virtually any nucleic acid molecule, including the isolated oligomers of the present invention.

**[0193]** As detailed below, the pharmaceutical compositions of the present invention may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (3) topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually; (6) ocularly; (7) transdermally; or (8) nasally.

**[0194]** The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0195] The phrase “pharmaceutically-acceptable carrier” as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler,

diluent, excipient, manufacturing aid (e.g., lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient.

**[0196]** Some examples of materials that can serve as pharmaceutically-acceptable carriers include, without limitation: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

**[0197]** Additional non-limiting examples of agents suitable for formulation with the antisense oligomers of the instant invention include: PEG conjugated nucleic acids, phospholipid conjugated nucleic acids, nucleic acids containing lipophilic moieties, phosphorothioates, P-glycoprotein inhibitors (such as Pluronic P85) which can enhance entry of drugs into various tissues; biodegradable polymers, such as poly (DL-lactide-coglycolide) microspheres for sustained release delivery after implantation (Emerich, D F et al., 1999, *Cell Transplant*, 8, 47-58) Alkermes, Inc. Cambridge, Mass.; and loaded nanoparticles, such as those made of polybutylcyanoacrylate, which can deliver drugs across the blood brain barrier and can alter neuronal uptake mechanisms (*Prog Neuropsychopharmacol Biol Psychiatry*, 23, 941-949, 1999).

**[0198]** The invention also features the use of the composition comprising surface-modified liposomes containing poly (ethylene glycol) lipids (PEG-modified, branched and unbranched or combinations thereof, or long-circulating liposomes or stealth liposomes). Oligomers of the invention can also comprise covalently attached PEG molecules of various



US 2010/0130591 A1

May 27, 2010

16

molecular weights. These formulations offer a method for increasing the accumulation of drugs in target tissues. This class of drug carriers resists opsonization and elimination by the mononuclear phagocytic system (MPS or RES), thereby enabling longer blood circulation times and enhanced tissue exposure for the encapsulated drug (Lasic et al. *Chem. Rev.* 1995, 95, 2601-2627; Ishiwata et al., *Chem. Pharm. Bull.* 1995, 43, 1005-1011). Such liposomes have been shown to accumulate selectively in tumors, presumably by extravasation and capture in the neovascularized target tissues (Lasic et al., *Science* 1995, 267, 1275-1276; Oku et al., 1995, *Biochim. Biophys. Acta*, 1238, 86-90). The long-circulating liposomes enhance the pharmacokinetics and pharmacodynamics of DNA and RNA, particularly compared to conventional cationic liposomes which are known to accumulate in tissues of the MPS (Liu et al., *J. Biol. Chem.* 1995, 42, 24864-24870; Choi et al., International PCT Publication No. WO 96/10391; Ansell et al., International PCT Publication No. WO 96/10390; Holland et al., International PCT Publication No. WO 96/10392). Long-circulating liposomes are also likely to protect drugs from nuclease degradation to a greater extent compared to cationic liposomes, based on their ability to avoid accumulation in metabolically aggressive MPS tissues such as the liver and spleen.

**[0199]** In a further embodiment, the present invention includes oligomer compositions prepared for delivery as described in U.S. Pat. Nos. 6,692,911, 7,163,695 and 7,070,807. In this regard, in one embodiment, the present invention provides an oligomer of the present invention in a composition comprising copolymers of lysine and histidine (HK) as described in U.S. Pat. Nos. 7,163,695, 7,070,807, and 6,692,911 either alone or in combination with PEG (e.g., branched or unbranched PEG or a mixture of both), in combination with PEG and a targeting moiety or any of the foregoing in combination with a crosslinking agent. In certain embodiments, the present invention provides antisense oligomers in compositions comprising gluconic-acid-modified polyhistidine or gluconylated-polyhistidine/transferrin-polylysine. One skilled in the art will also recognize that amino acids with properties similar to His and Lys may be substituted within the composition.

**[0200]** Certain embodiments of the oligomers described herein may contain a basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable acids. The term "pharmaceutically-acceptable salts" in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared in situ in the administration vehicle or the dosage form manufacturing process, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed during subsequent purification. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, e.g., Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19).

**[0201]** The pharmaceutically acceptable salts of the subject oligomers include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from non-toxic organic or inorganic acids. For example, such conventional

nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like.

**[0202]** In certain embodiments, the oligomers of the present invention may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can likewise be prepared in situ in the administration vehicle or the dosage form manufacturing process, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. (See, e.g., Berge et al., *supra*).

**[0203]** Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

**[0204]** Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

**[0205]** Formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred percent, this amount will range from about 0.1 percent to about ninety-nine percent of active ingredient, preferably from about 5 percent to about 70 percent, most preferably from about 10 percent to about 30 percent.

**[0206]** In certain embodiments, a formulation of the present invention comprises an excipient selected from cyclodextrins, celluloses, liposomes, micelle forming agents, e.g., bile acids, and polymeric carriers, e.g., polyesters and poly-

US 2010/0130591 A1

May 27, 2010

17

anhydrides; and an oligomer of the present invention. In certain embodiments, an aforementioned formulation renders orally bioavailable an oligomer of the present invention.

**[0207]** Methods of preparing these formulations or compositions include the step of bringing into association an oligomer of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

**[0208]** Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. An oligomer of the present invention may also be administered as a bolus, electuary or paste.

**[0209]** In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules, trouches and the like), the active ingredient may be mixed with one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds and surfactants, such as poloxamer and sodium lauryl sulfate; (7) wetting agents, such as, for example, cetyl alcohol, glycerol monostearate, and non-ionic surfactants; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, zinc stearate, sodium stearate, stearic acid, and mixtures thereof; (10) coloring agents; and (11) controlled release agents such as crospovidone or ethyl cellulose. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

**[0210]** A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (e.g., gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

**[0211]** The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric

coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

**[0212]** Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

**[0213]** Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

**[0214]** Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

**[0215]** Formulations for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

**[0216]** Formulations or dosage forms for the topical or transdermal administration of an oligomer as provided herein include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active oligomers may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any preservatives, buffers, or propellants which may be required. The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

**[0217]** Powders and sprays can contain, in addition to an oligomer of the present invention, excipients such as lactose,

US 2010/0130591 A1

May 27, 2010

18

talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

[0218] Transdermal patches have the added advantage of providing controlled delivery of an oligomer of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the oligomer in the proper medium. Absorption enhancers can also be used to increase the flux of the agent across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the agent in a polymer matrix or gel, among other methods known in the art.

[0219] Pharmaceutical compositions suitable for parenteral administration may comprise one or more oligomers of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain sugars, alcohols, antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents. Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0220] These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms upon the subject oligomers may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0221] In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility, among other methods known in the art. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

[0222] Injectable depot forms may be made by forming microencapsule matrices of the subject oligomers in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of oligomer to polymer, and the nature of the particular polymer employed, the rate of oligomer release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides).

Depot injectable formulations may also be prepared by entrapping the drug in liposomes or microemulsions that are compatible with body tissues.

[0223] When the oligomers of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99% (more preferably, 10 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

[0224] As noted above, the formulations or preparations of the present invention may be given orally, parenterally, topically, or rectally. They are typically given in forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories.

[0225] The phrases "parenteral administration" and "administered parenterally" as used herein mean modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0226] The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

[0227] Regardless of the route of administration selected, the oligomers of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, may be formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art. Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being unacceptably toxic to the patient.

[0228] The selected dosage level will depend upon a variety of factors including the activity of the particular oligomer of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion or metabolism of the particular oligomer being employed, the rate and extent of absorption, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular oligomer employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0229] A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a



US 2010/0130591 A1

May 27, 2010

19

suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, oral, intravenous, intracerebroventricular and subcutaneous doses of the compounds of this invention for a patient, when used for the indicated effects, will range from about 0.0001 to about 100 mg per kilogram of body weight per day.

**[0230]** If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. In certain situations, dosing is one administration per day. In certain embodiments, dosing is one or more administration per every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 days, or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 weeks, or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 months, as needed, to maintain the desired expression of a functional dystrophin protein.

**[0231]** Nucleic acid molecules can be administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres, as described herein and known in the art. In certain embodiments, microemulsification technology may be utilized to improve bioavailability of lipophilic (water insoluble) pharmaceutical agents. Examples include Trimetrine (Dordunoo, S. K., et al., *Drug Development and Industrial Pharmacy*, 17(12), 1685-1713, 1991 and REV 5901 (Sheen, P. C., et al., *J Pharm Sci* 80(7), 712-714, 1991). Among other benefits, microemulsification provides enhanced bioavailability by preferentially directing absorption to the lymphatic system instead of the circulatory system, which thereby bypasses the liver, and prevents destruction of the compounds in the hepatobiliary circulation.

**[0232]** In one aspect of invention, the formulations contain micelles formed from an oligomer as provided herein and at least one amphiphilic carrier, in which the micelles have an average diameter of less than about 100 nm. More preferred embodiments provide micelles having an average diameter less than about 50 nm, and even more preferred embodiments provide micelles having an average diameter less than about 30 nm, or even less than about 20 nm.

**[0233]** While all suitable amphiphilic carriers are contemplated, the presently preferred carriers are generally those that have Generally-Recognized-as-Safe (GRAS) status, and that can both solubilize the compound of the present invention and microemulsify it at a later stage when the solution comes into a contact with a complex water phase (such as one found in human gastro-intestinal tract). Usually, amphiphilic ingredients that satisfy these requirements have HLB (hydrophilic to lipophilic balance) values of 2-20, and their structures contain straight chain aliphatic radicals in the range of C-6 to C-20. Examples are polyethylene-glycolized fatty glycerides and polyethylene glycols.

**[0234]** Examples of amphiphilic carriers include saturated and monounsaturated polyethyleneglycolized fatty acid glycerides, such as those obtained from fully or partially hydrogenated various vegetable oils. Such oils may advantageously consist of tri-, di-, and mono-fatty acid glycerides and di- and mono-polyethyleneglycol esters of the corresponding fatty acids, with a particularly preferred fatty acid composition including capric acid 4-10, capric acid 3-9, lauric acid 40-50, myristic acid 14-24, palmitic acid 4-14 and stearic acid

5-15%. Another useful class of amphiphilic carriers includes partially esterified sorbitan and/or sorbitol, with saturated or mono-unsaturated fatty acids (SPAN-series) or corresponding ethoxylated analogs (TWEEN-series).

**[0235]** Commercially available amphiphilic carriers may be particularly useful, including Gelucire-series, Labrafil, Labrasol, or Lauroglycol (all manufactured and distributed by Gattefossé Corporation, Saint Priest, France), PEG-mono-oleate, PEG-di-oleate, PEG-mono-laurate and di-laurate, Lecithin, Polysorbate 80, etc (produced and distributed by a number of companies in USA and worldwide).

**[0236]** In certain embodiments, the delivery may occur by use of liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, for the introduction of the compositions of the present invention into suitable host cells. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, a nanoparticle or the like. The formulation and use of such delivery vehicles can be carried out using known and conventional techniques.

**[0237]** Hydrophilic polymers suitable for use in the present invention are those which are readily water-soluble, can be covalently attached to a vesicle-forming lipid, and which are tolerated in vivo without toxic effects (i.e., are biocompatible). Suitable polymers include polyethylene glycol (PEG), polylactic (also termed polylactide), polyglycolic acid (also termed polyglycolide), a polylactic-polyglycolic acid copolymer, and polyvinyl alcohol. In certain embodiments, polymers have a molecular weight of from about 100 or 120 daltons up to about 5,000 or 10,000 daltons, or from about 300 daltons to about 5,000 daltons. In other embodiments, the polymer is polyethyleneglycol having a molecular weight of from about 100 to about 5,000 daltons, or having a molecular weight of from about 300 to about 5,000 daltons. In certain embodiments, the polymer is polyethyleneglycol of 750 daltons (PEG(750)). Polymers may also be defined by the number of monomers therein; a preferred embodiment of the present invention utilizes polymers of at least about three monomers, such PEG polymers consisting of three monomers (approximately 150 daltons).

**[0238]** Other hydrophilic polymers which may be suitable for use in the present invention include polyvinylpyrrolidone, polymethoxazoline, polyethyloxazoline, polyhydroxypropyl methacrylamide, polymethacrylamide, polydimethylacrylamide, and derivatized celluloses such as hydroxymethylcellulose or hydroxyethylcellulose.

**[0239]** In certain embodiments, a formulation of the present invention comprises a biocompatible polymer selected from the group consisting of polyamides, polycarbonates, polyalkylenes, polymers of acrylic and methacrylic esters, polyvinyl polymers, polyglycolides, polysiloxanes, polyurethanes and co-polymers thereof, celluloses, polypropylene, polyethylenes, polystyrene, polymers of lactic acid and glycolic acid, polyanhydrides, poly(ortho)esters, poly(butic acid), poly(valeric acid), poly(lactide-co-caprolactone), polysaccharides, proteins, polyhyaluronic acids, polycyanoacrylates, and blends, mixtures, or copolymers thereof.

**[0240]** Cyclodextrins are cyclic oligosaccharides, consisting of 6, 7 or 8 glucose units, designated by the Greek letter  $\alpha$ ,  $\beta$ , or  $\gamma$ , respectively. The glucose units are linked by  $\alpha$ -1,4-glucosidic bonds. As a consequence of the chair conformation of the sugar units, all secondary hydroxyl groups (at C-2, C-3) are located on one side of the ring, while all the primary

US 2010/0130591 A1

May 27, 2010

20

hydroxyl groups at C-6 are situated on the other side. As a result, the external faces are hydrophilic, making the cyclodextrins water-soluble. In contrast, the cavities of the cyclodextrins are hydrophobic, since they are lined by the hydrogen of atoms C-3 and C-5, and by ether-like oxygens. These matrices allow complexation with a variety of relatively hydrophobic compounds, including, for instance, steroid compounds such as  $17\alpha$ -estradiol (see, e.g., van Uden et al. *Plant Cell Tiss. Org. Cult.* 38:1-3-113 (1994)). The complexation takes place by Van der Waals interactions and by hydrogen bond formation. For a general review of the chemistry of cyclodextrins, see, Wenz, *Agnew. Chem. Int. Ed. Engl.*, 33:803-822 (1994).

[0241] The physico-chemical properties of the cyclodextrin derivatives depend strongly on the kind and the degree of substitution. For example, their solubility in water ranges from insoluble (e.g., triacetyl-beta-cyclodextrin) to 147% soluble (w/v) (G-2-beta-cyclodextrin). In addition, they are soluble in many organic solvents. The properties of the cyclodextrins enable the control over solubility of various formulation components by increasing or decreasing their solubility.

[0242] Numerous cyclodextrins and methods for their preparation have been described. For example, Parmeter (I), et al. (U.S. Pat. No. 3,453,259) and Gramera, et al. (U.S. Pat. No. 3,459,731) described electroneutral cyclodextrins. Other derivatives include cyclodextrins with cationic properties [Parmeter (II), U.S. Pat. No. 3,453,257], insoluble crosslinked cyclodextrins (Solms, U.S. Pat. No. 3,420,788), and cyclodextrins with anionic properties [Parmeter (III), U.S. Pat. No. 3,426,011]. Among the cyclodextrin derivatives with anionic properties, carboxylic acids, phosphorous acids, phosphinous acids, phosphonic acids, phosphoric acids, thiophosphonic acids, thiosulphinic acids, and sulfonic acids have been appended to the parent cyclodextrin [see, Parmeter (III), *supra*]. Furthermore, sulfoalkyl ether cyclodextrin derivatives have been described by Stella, et al. (U.S. Pat. No. 5,134,127).

[0243] Liposomes consist of at least one lipid bilayer membrane enclosing an aqueous internal compartment. Liposomes may be characterized by membrane type and by size. Small unilamellar vesicles (SUVs) have a single membrane and typically range between 0.02 and 0.05  $\mu\text{m}$  in diameter; large unilamellar vesicles (LUVs) are typically larger than 0.05  $\mu\text{m}$ . Oligolamellar large vesicles and multilamellar vesicles have multiple, usually concentric, membrane layers and are typically larger than 0.1  $\mu\text{m}$ . Liposomes with several nonconcentric membranes, i.e., several smaller vesicles contained within a larger vesicle, are termed multivesicular vesicles.

[0244] One aspect of the present invention relates to formulations comprising liposomes containing an oligomer of the present invention, where the liposome membrane is formulated to provide a liposome with increased carrying capacity. Alternatively or in addition, the compound of the present invention may be contained within, or adsorbed onto, the liposome bilayer of the liposome. An oligomer of the present invention may be aggregated with a lipid surfactant and carried within the liposome's internal space; in these cases, the liposome membrane is formulated to resist the disruptive effects of the active agent-surfactant aggregate.

[0245] According to one embodiment of the present invention, the lipid bilayer of a liposome contains lipids derivatized with polyethylene glycol (PEG), such that the PEG chains

extend from the inner surface of the lipid bilayer into the interior space encapsulated by the liposome, and extend from the exterior of the lipid bilayer into the surrounding environment.

[0246] Active agents contained within liposomes of the present invention are in solubilized form. Aggregates of surfactant and active agent (such as emulsions or micelles containing the active agent of interest) may be entrapped within the interior space of liposomes according to the present invention. A surfactant acts to disperse and solubilize the active agent, and may be selected from any suitable aliphatic, cycloaliphatic or aromatic surfactant, including but not limited to biocompatible lysophosphatidylcholines (LPGs) of varying chain lengths (for example, from about C14 to about C20). Polymer-derivatized lipids such as PEG-lipids may also be utilized for micelle formation as they will act to inhibit micelle/membrane fusion, and as the addition of a polymer to surfactant molecules decreases the CMC of the surfactant and aids in micelle formation. Preferred are surfactants with CMOs in the micromolar range; higher CMC surfactants may be utilized to prepare micelles entrapped within liposomes of the present invention.

[0247] Liposomes according to the present invention may be prepared by any of a variety of techniques that are known in the art. See, e.g., U.S. Pat. No. 4,235,871; Published PCT applications WO 96/14057; New RRC, *Liposomes: A practical approach*, IRL Press, Oxford (1990), pages 33-104; Lasic DD, *Liposomes from physics to applications*, Elsevier Science Publishers BV, Amsterdam, 1993. For example, liposomes of the present invention may be prepared by diffusing a lipid derivatized with a hydrophilic polymer into preformed liposomes, such as by exposing preformed liposomes to micelles composed of lipid-grafted polymers, at lipid concentrations corresponding to the final mole percent of derivatized lipid which is desired in the liposome. Liposomes containing a hydrophilic polymer can also be formed by homogenization, lipid-field hydration, or extrusion techniques, as are known in the art.

[0248] In another exemplary formulation procedure, the active agent is first dispersed by sonication in a lysophosphatidylcholine or other low CMC surfactant (including polymer grafted lipids) that readily solubilizes hydrophobic molecules. The resulting micellar suspension of active agent is then used to rehydrate a dried lipid sample that contains a suitable mole percent of polymer-grafted lipid, or cholesterol. The lipid and active agent suspension is then formed into liposomes using extrusion techniques as are known in the art, and the resulting liposomes separated from the unencapsulated solution by standard column separation.

[0249] In one aspect of the present invention, the liposomes are prepared to have substantially homogeneous sizes in a selected size range. One effective sizing method involves extruding an aqueous suspension of the liposomes through a series of polycarbonate membranes having a selected uniform pore size; the pore size of the membrane will correspond roughly with the largest sizes of liposomes produced by extrusion through that membrane. See e.g., U.S. Pat. No. 4,737,323 (Apr. 12, 1988). In certain embodiments, reagents such as DharmaFECT® and Lipofectamine® may be utilized to introduce polynucleotides or proteins into cells.

[0250] The release characteristics of a formulation of the present invention depend on the encapsulating material, the concentration of encapsulated drug, and the presence of release modifiers. For example, release can be manipulated to

US 2010/0130591 A1

May 27, 2010

21

be pH dependent, for example, using a pH sensitive coating that releases only at a low pH, as in the stomach, or a higher pH, as in the intestine. An enteric coating can be used to prevent release from occurring until after passage through the stomach. Multiple coatings or mixtures of cyanamide encapsulated in different materials can be used to obtain an initial release in the stomach, followed by later release in the intestine. Release can also be manipulated by inclusion of salts or pore forming agents, which can increase water uptake or release of drug by diffusion from the capsule. Excipients which modify the solubility of the drug can also be used to control the release rate. Agents which enhance degradation of the matrix or release from the matrix can also be incorporated. They can be added to the drug, added as a separate phase (i.e., as particulates), or can be co-dissolved in the polymer phase depending on the compound. In most cases the amount should be between 0.1 and thirty percent (w/w polymer). Types of degradation enhancers include inorganic salts such as ammonium sulfate and ammonium chloride, organic acids such as citric acid, benzoic acid, and ascorbic acid, inorganic bases such as sodium carbonate, potassium carbonate, calcium carbonate, zinc carbonate, and zinc hydroxide, and organic bases such as protamine sulfate, spermine, choline, ethanolamine, diethanolamine, and triethanolamine and surfactants such as Tween® and Pluronic®. Pore forming agents which add microstructure to the matrices (i.e., water soluble compounds such as inorganic salts and sugars) are added as particulates. The range is typically between one and thirty percent (w/w polymer).

[0251] Uptake can also be manipulated by altering residence time of the particles in the gut. This can be achieved, for example, by coating the particle with, or selecting as the encapsulating material, a mucosal adhesive polymer. Examples include most polymers with free carboxyl groups, such as chitosan, celluloses, and especially polyacrylates (as used herein, polyacrylates refers to polymers including acrylate groups and modified acrylate groups such as cyanoacrylates and methacrylates).

[0252] An oligomer may be formulated to be contained within, or, adapted to release by a surgical or medical device or implant. In certain aspects, an implant may be coated or otherwise treated with an oligomer. For example, hydrogels, or other polymers, such as biocompatible and/or biodegradable polymers, may be used to coat an implant with the compositions of the present invention (i.e., the composition may be adapted for use with a medical device by using a hydrogel or other polymer). Polymers and copolymers for coating medical devices with an agent are well-known in the art. Examples of implants include, but are not limited to, stents, drug-eluting stents, sutures, prosthesis, vascular catheters, dialysis catheters, vascular grafts, prosthetic heart valves, cardiac pacemakers, implantable cardioverter defibrillators, IV needles, devices for bone setting and formation, such as pins, screws, plates, and other devices, and artificial tissue matrices for wound healing.

[0253] In addition to the methods provided herein, the oligomers for use according to the invention may be formulated for administration in any convenient way for use in human or veterinary medicine, by analogy with other pharmaceuticals. The antisense oligomers and their corresponding formulations may be administered alone or in combination with other

therapeutic strategies in the treatment of muscular dystrophy, such as myoblast transplantation, stem cell therapies, administration of aminoglycoside antibiotics, proteasome inhibitors, and up-regulation therapies (e.g., upregulation of utrophin, an autosomal paralogue of dystrophin).

[0254] All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

[0255] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims. The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

#### REFERENCES

- [0256] Aartsma-Rus, A., A. A. Janson, et al. (2004). "Antisense-induced multiexon skipping for Duchenne muscular dystrophy makes more sense." *Am J Hum Genet.* 74(1): 83-92.
- [0257] Dunckley, M. G., I. C. Eperon, et al. (1997). "Modulation of splicing in the DMD gene by antisense oligoribonucleotides." *Nucleosides & Nucleotides* 16(7-9): 1665-1668.
- [0258] Dunckley, M. G., M. Manoharan, et al. (1998). "Modification of splicing in the dystrophin gene in cultured Mdx muscle cells by antisense oligoribonucleotides." *Hum Mol Genet.* 7(7): 1083-90.
- [0259] Errington, S. J., C. J. Mann, et al. (2003). "Target selection for antisense oligonucleotide induced exon skipping in the dystrophin gene." *J Gene Med* 5(6): 518-27.
- [0260] Jearawiriyapaisarn, N., H. M. Moulton, et al. (2008). "Sustained Dystrophin Expression Induced by Peptide-conjugated Morpholino Oligomers in the Muscles of mdx Mice." *Mol. Ther.*
- [0261] Lu, Q. L., C. J. Mann, et al. (2003). "Functional amounts of dystrophin produced by skipping the mutated exon in the mdx dystrophic mouse." *Nat Med* 9(8): 1009-14.
- [0262] Mann, C. J., K. Honeyman, et al. (2002). "Improved antisense oligonucleotide induced exon skipping in the mdx mouse model of muscular dystrophy." *J Gene Med* 4(6): 644-54.
- [0263] Marshall, N. B., S. K. Oda, et al. (2007). "Arginine-rich cell-penetrating peptides facilitate delivery of antisense oligomers into murine leukocytes and alter pre-mRNA splicing." *Journal of Immunological Methods* 325 (1-2): 114-126.
- [0264] Matsuo, M., T. Masumura, et al. (1991). "Exon skipping during splicing of dystrophin mRNA precursor due to an intraexon deletion in the dystrophin gene of Duchenne muscular dystrophy kobe." *J Clin Invest* 87(6): 2127-31.



US 2010/0130591 A1

May 27, 2010

22

- [0265] Monaco, A. P., C. J. Bertelson, et al. (1988). "An explanation for the phenotypic differences between patients bearing partial deletions of the DMD locus." *Genomics* 2(1): 90-5.
- [0266] Pramono, Z. A., Y. Takeshima, et al. (1996). "Induction of exon skipping of the dystrophin transcript in lymphoblastoid cells by transfecting an antisense oligodeoxynucleotide complementary to an exon recognition sequence." *Biochem Biophys Res Commun* 226(2): 445-9.
- [0267] Sazani, P., R. Kole, et al. (2007). Splice switching oligomers for the TNF superfamily receptors and their use in treatment of disease. PCT WO2007058894, University of North Carolina
- [0268] Sierakowska, H., M. J. Sambade, et al. (1996). "Repair of thalassemic human beta-globin mRNA in mammalian cells by antisense oligonucleotides." *Proc Natl Acad Sci USA* 93(23): 12840-4.
- [0269] Summerton, J. and D. Weller (1997). "Morpholino antisense oligomers: design, preparation, and properties." *Antisense Nucleic Acid Drug Dev* 7(3): 187-95.
- [0270] Takeshima, Y., H. Nishio, et al. (1995). "Modulation of in vitro splicing of the upstream intron by modifying an intra-exon sequence which is deleted from the dystrophin gene in dystrophin Kobe." *J Clin Invest* 95(2): 515-20.
- [0271] van Deutekom, J. C., M. Bremmer-Bout, et al. (2001). "Antisense-induced exon skipping restores dystrophin expression in DMD patient derived muscle cells." *Hum Mol Genet.* 10(15): 1547-54.
- [0272] van Deutekom, J. C., A. A. Janson, et al. (2007). "Local dystrophin restoration with antisense oligonucleotide PRO051." *N Engl J Med* 357(26): 2677-86.
- [0273] Wilton, S. D., A. M. Fall, et al. (2007). "Antisense oligonucleotide-induced exon skipping across the human dystrophin gene transcript." *Mol Ther* 15(7): 1288-96.
- [0274] Wilton, S. D., F. Lloyd, et al. (1999). "Specific removal of the nonsense mutation from the mdx dystrophin mRNA using antisense oligonucleotides." *Neuromuscul Disord* 9(5): 330-8.
- [0275] Wu, B., H. M. Moulton, et al. (2008). "Effective rescue of dystrophin improves cardiac function in dystrophin-deficient mice by a modified morpholino oligomer." *Proc Natl Acad Sci USA* 105(39): 14814-9.
- [0276] Yin, H., H. M. Moulton, et al. (2008). "Cell-penetrating peptide-conjugated antisense oligonucleotides restore systemic muscle and cardiac dystrophin expression and function." *Hum Mol Genet.* 17(24): 3909-18.
- [0279] Media was aspirated from T75 flasks; cells were washed once in warmed PBS and aspirated. 3 mL of Trypsin/EDTA, warmed in a 37° C. water bath, was added to each T75. Cells were incubated at 37° C. 5-5 minutes until, with gentle agitation, they released from the flask. Cell suspension was transferred to a 15.0 mL conical tube; flasks were rinsed with 1.0 mL of Trypsin/EDTA solution to gather remaining cells. Cells were counted with a Vi-Cell XR cell counter (Beckman Coulter). Cells were seeded into tissue culture-treated 12-well plates (Falcon) at 2.0×10<sup>5</sup> viable cells per well in 1.0 mL media. Cells were incubated overnight in a 37° C. incubator at 5.0% CO<sub>2</sub>.
- [0280] Twelve-well seeded plates were examined for even cellular distribution and plate adherence. Lyophilized peptide conjugated phosphorodiamidate morpholino oligomers (PPMOs) were re-suspended at 2.0 mM in nuclease-free water (Ambion), and kept on ice during cell treatment; to verify molarity, PPMOs were measured using a Nanoprop 2000 spectrophotometer (Thermo Scientific). Immediately prior to PPMO treatment, media was aspirated, and cells were rinsed in warmed PBS. PPMOs were diluted in warmed media to the desired molarity; cells were treated in a total of 1.0 mL PPMO per well. PPMOs were tested in triplicate. For no-treatment controls, fresh, warmed media was added in 1.0 mL total volume. Cells were incubated for 48 hours in a 37° C. incubator at 5.0% CO<sub>2</sub>.
- [0281] RNA Extraction
- [0282] Media was aspirated, and cells were rinsed in warmed PBS. RNA was extracted with the QuickGene-Mini80 system, QuickGene RNA cultured cell HC kit S, and MagNalyser with ceramic bead homogenization using the manufacturers' recommended protocols. Briefly, cells were lysed in treatment plates with 350 uL LRP (10 uL β-Mercaptoethanol added per 100 uL LRP) lysis buffer; homogenate was gently triturated to ensure full lysis, and transferred to MagNalyser tubes. Tubes were spun at 2800 rpm for 30 seconds in the MagNalyser to ensure full homogenization, and iced briefly. 50 uL SRP solubilization buffer was added and homogenate was vortexed for 15 seconds. 170 uL >99% ethanol was added to each tube, and homogenate was vortexed for 60 seconds. Homogenate was flash-spun and transferred to Mini80 RNA cartridges, samples were pressurized and flow-through was discarded. Cartridges were washed in 750 uL WRP wash buffer and pressurized. 40 uL of DNase solution (1.25 uL Qiagen DNaseI, 35 uL RDD Buffer, 3.75 uL nuclease-free water) was added directly to the cartridge membrane; cartridges were incubated four minutes at room temperature. Cartridges were washed twice with 750 uL WRP, pressurizing after each wash. Cartridges were placed over nuclease-free tubes. 50 uL CRP elution buffer was added to each membrane; membranes were incubated for five minutes at room-temperature. Cartridges were pressurized and eluate was collected. RNA was stored at -80° C. pending quantification. RNA was quantified using the Nanoprop™ 2000 spectrophotometer.

## EXAMPLES

### Materials and Methods

- [0277] Cells and Tissue Culture Treatment Conditions
- [0278] Human Rhabdomyosarcoma cells (ATCC, CCL-136; RD cells) preserved in a 5% DMSO solution (Sigma) at a low passage number were thawed in a 37° C. water bath until the ice sliver was no longer visible. Cells were seeded into tissue culture-treated T75 flasks (Nunc) at 1.5×10<sup>6</sup> cells/flask in 24 mL of warmed DMEM with L-Glutamine (HyClone), 10% fetal bovine serum, and 1% Penicillin-Streptomycin antibiotic solution (CelGro); after 24 hours, media was aspirated, cells were washed once in warmed PBS, and fresh media was added. Cells were grown to 80% confluence in a 37° C. incubator at 5.0% CO<sub>2</sub>.
- [0283] Nested RT-PCR
- [0284] Primer-specific, exon-specific, optimized nested RT-PCR amplification was performed using the primer pair sets for each dystrophin exon as shown below in Table 1.

US 2010/0130591 A1

23

May 27, 2010

TABLE 1

Primer pair sets used to PCR amplify human dystrophin mRNA to detect exon-skipping.					
Name	F/R	I/O	Sequence (5'-3')	Exon Purpose	SEQ ID NO:
PS170	F	O	CCAGAGCTTTACCTGAGAAACAAG	48 Detection	640
PS172	F	I	CCAGCCACTCAGCCAGTGAAG	49 of Exon 50	641
PS174	R	I	CGATCCGTAATGATTGTTCTAGCC	52 and 51	642
PS176	R	O	CATTTCAATCAACTGTTGCCTCCG	53 Skipping in Human Dystrophin	643
PS186	F	O	CAATGCTCCTGACCTCTGTGC	42 Detection	644
PS187	F	I	GTCTACAACAAGCTCAGGTCG	43 of Exon 44	645
PS189	F	I	GCAATGTTATCTGCTTCCTCCAACC	46 and 45	646
PS190	R	O	GCTCTTTCCAGGTTCAAGTGG	46 Skipping in Human Dystrophin	647
PS192	F	O	CTTGGACAGAACTTACCGACTGG	51 Detection	648
PS193	F	I	GCAGGATTTGGAACAGAGGCG	52 of Exon 53	649
PS195	R	I	CATCTACATTTGCTGCCACTGG	54 Skipping in	650
PS197	R	O	GTTTCTTCCAAGCAGCCTCTCG	55 Human Dystrophin	651

**[0285]** The indicated primer pairs are shown as either forward or reverse (F/R) and either outer or inner primer pairs (I/O) corresponding to primary or secondary amplifications, respectively. The location of the primer target is indicated in the Exon column and the Purpose indicates the exon-skipping events can be detected. For example, PS170 and PS176 primers amplify a region from exon 48 to 53 in the primary amplification. Primers PS172 and PS174 then amplify a region from exon 49 to 52 in the secondary amplification. This nested PCR reaction will detect exon skipping of both exons 50 and/or exon 51. The specific nested RT-PCR reaction conditions are provided below.

**[0286]** RNA extracted from treated cells (described above) was diluted to 20 ng/ul for all samples.

TABLE 2

Reaction setup for RT-PCR and primary amplification (50 µl reaction):	
2x Reaction mix	25 µl
PS XXX Forward Primer (30 µM)	0.5 µl
(see Table 1)	
PS XXX Reverse Primer (30 µM)	0.5 µl
(see Table 1)	
Superscript III Platinum Taq mix	2 µl
Template RNA (20 ng/ul)	10 µl
Nuclease-Free Water (50 µl total volume)	12 µl

TABLE 3

RT-PCR and primary amplification program:			
	Temperature	Time	
Reverse Transcription	55° C.	30 minutes	
RT Inactivation	94° C.	2 minutes	
Denaturing	94° C.	1 minute	8 Cycles
Annealing	59° C.	1 minute	

TABLE 3-continued

RT-PCR and primary amplification program:		
	Temperature	Time
Extension	68° C.	1 minute
	4° C.	∞

TABLE 4

Reaction setup for nested secondary amplification (50 µl reaction):	
10x PCR Buffer	5 µl
dNTP solution (10 mM)	0.5 µl
50 mM MgCl	1.5 µl
PS XXX Forward Primer (30 µM)	0.33 µl
(see Table 1)	
PS XXX Reverse Primer (30 µM)	0.33 µl
(see Table 1)	
Platinum Taq DNA polymerase	0.2 µl
0.1 mM Cy5-dCTP	1 µl
RT-PCR product (from Step 1)	1 µl
Nuclease-Free Water (50 µl total volume)	40.15 µl

TABLE 5

Nested secondary amplification program:			
	Temperature	Time	
Primary Denature	94° C.	3 minutes	
Denaturing	94° C.	45 seconds	28-30 Cycles
Annealing	59° C.	30 seconds	
Extension	68° C.	1 minute	
	4° C.	∞	



US 2010/0130591 A1

May 27, 2010

24

**[0287]** Gel Electrophoresis Analysis

**[0288]** Ten microliters of 5× Ficoll loading dye was added to each 50 microliter nested RT-PCR reaction. Fifteen microliters of PCR/dye mixture was run on a 10% TBE gel at 300 volts for 30 minutes. After electrophoresis, the gel was washed in diH<sub>2</sub>O for at least one hour, changing the water every 30 minutes. The gel was then scanned on a Typhoon Trio Variable Mode Imager (GE Healthcare). For exon 44 skipping, the nested RT-PCR product from full-length dystrophin transcript is 571 bp, and 423 bp from Exon 44-skipped mRNA (exon 44 is 148 bp). For exon 45, the nested RT-PCR product from full-length dystrophin transcript is 571 bp, and 395 bp from Exon 45-skipped mRNA (exon 45 is 176 bp). For exon 53, the PCR product from full-length dystrophin transcript is 365 bp, and 153 bp from exon 53-skipped mRNA (exon 53 is 212 bp).

**[0289]** The gel images were subjected to quantitative analysis by measuring the band intensities of the full-length PCR product compared to the exon-skipped product. In some cases, the percent skipping at a fixed PPMO concentration (e.g., 3 micromolar) was used to determine the relative activity of a series of PPMO to induce exon skipping of a given exon. In other situations, a PPMO dose-range was used to treat cells (e.g., 0.1, 0.3, 1.0, 3.0 and 10 micromolar) and an EC<sub>50</sub> was calculated based on the percent skipping induced at each concentration.

## Example 1

## Exon 51 Scan

**[0290]** A series of overlapping antisense PPMOs that target human dystrophin exon 51 were designed, synthesized and used to treat either human rhabdomyosarcoma cells (RD cells) or primary human skeletal muscle cells. This strategy is termed an “exon scan” and was used similarly for several other dystrophin exons as described below. All the PPMOs were synthesized as peptide-conjugated PMO (PPMO) using the CP06062 peptide (SEQ ID NO: 578) and a 3' terminal PMO linkage. For exon 51, a series of 26 PPMOs, each 26 bases in length, were made (SEQ ID NOS: 309-311, 314, 316, 317, 319, 321, 323, 324, 326, 327, 329-331, 333, 335, 336, 338-345) as shown in FIG. 2A. The PPMOs were evaluated for exon skipping efficacy by treating RD cells at various concentrations as described above in the Materials and Methods. Three PPMOs (SEQ ID NOS: 324, 326 and 327) were identified as effective in inducing exon-skipping and selected for additional evaluation. Dose-ranging experiments in RD cells and primary human skeletal muscle cells were used to confirm the relative efficacy of these three PPMO sequences. SEQ ID NO: 327 was shown to be most effective at inducing exon 51 skipping as shown in FIGS. 2B and 2C.

**[0291]** A comparison of the relative effectiveness of SEQ ID NO: 327 to other exon 51-targeted antisense sequences was performed in RD cells and primary human skeletal muscle cells, as described above. All the evaluated sequences were made as peptide-conjugated PMOs using the CP06062 peptide (SEQ ID NO: 578). This allowed direct comparison of the relative effectiveness of the antisense sequences without regard to antisense chemistry or cell delivery. The relative location of the certain exon 51-targeted oligos compared to SEQ ID NO: 327 is shown in FIG. 2D. As shown in FIG. 2C, there is a ranked hierarchy of exon-skipping effectiveness,

with SEQ ID NO: 327 being the most effective by at least a factor of several-fold compared to other sequences.

## Example 2

## Exon 50 Scan

**[0292]** A series of overlapping antisense PPMOs that target human dystrophin exon 50 were designed and synthesized. For exon 50, a series of 17 PPMOs, each 25 bases in length, were made (SEQ ID NOS: 267, 269, 271, 273, 275, 277, 279, 280, 282 and 284-291) as shown in FIG. 3A. The PPMOs were evaluated for exon skipping efficacy by treating RD cells at various concentrations as described above in the Materials and Methods. Four PPMOs (SEQ ID NOS: 277, 287, 290 and 291) were identified as effective in inducing exon-skipping and selected for additional evaluation. Dose-ranging experiments in RD cells were used to confirm the relative efficacy of these four PMO sequences. SEQ ID NOS: 584 (AVI-5656) and 287 (AVI-5038) were shown to be most effective at inducing exon 50 skipping as shown in FIG. 3B. The EC<sub>50</sub> values were derived from the dose-ranging experiments and represent the calculated concentration where 50% of the PCR product is from the mRNA lacking exon 50 relative to the PCR product produced from the mRNA containing exon 50. Compared to other sequences (see, e.g., SEQ ID NOS: 584 and 585 correspond to SEQ ID NOS: 173 and 175 in WO2006/000057, respectively) AVI-5038 (SEQ ID NO: 287) is equivalent or better at inducing exon-skipping activity in the RD cell assay as shown in FIG. 3B.

## Example 3

## Exon 53 Scan

**[0293]** A series of overlapping antisense PPMOs that target human dystrophin exon 53 were designed and synthesized. For exon 53, a series of 24 PPMOs, each 25 bases in length, were made (SEQ ID NOS: 416, 418, 420, 422, 424, 426, 428, 429, 431, 433, 434, 436, 438-440 and 443-451) as shown in FIG. 4A. The PPMOs were evaluated for exon skipping efficacy by treating RD cells and primary human skeletal muscle cells at various concentrations as described above in the Materials and Methods. Three PPMOs (SEQ ID NOS: 428, 429 and 431) were identified as effective in inducing exon-skipping and selected for additional evaluation. Dose-ranging experiments in RD cells were used to confirm the relative efficacy of these three PMO sequences. SEQ ID NO: 429 was shown to be most effective at inducing exon 53 skipping as shown in FIGS. 4B-F. However, when compared to other exon 53 antisense sequences, SEQ ID NO: 429 proved identical to H53A(+23+47) which is listed as SEQ ID NO: 195 in WO2006/000057 and SEQ ID NO: 609 in the present application. Other sequences were compared to SEQ ID NO: 429 including H53A(+39+69) and H53A(-12+10) (listed as SEQ ID NOS: 193 and 199 in WO2006/000057, respectively) and h53AON1 (listed as SEQ ID NO: 39 in U.S. application Ser. No. 11/233,507) and listed as SEQ ID NOS: 608, 611 and 610, respectively, in the present application. All the evaluated sequences were made as peptide-conjugated PMOs using the CP06062 peptide (SEQ ID NO: 578). This allowed direct comparison of the relative effectiveness of the antisense sequences without regard to antisense chemistry or cell delivery.

US 2010/0130591 A1

May 27, 2010

25

ery. As shown in FIGS. 4I and 4G-H, SEQ ID NO: 429 was shown to be superior to each of these four sequences.

#### Example 4

##### Exon 44 Scan

[0294] A series of overlapping antisense PPMOs that target human dystrophin exon 44 were designed and synthesized. For exon 44, a series of PPMOs, each 25 bases in length, were made (SEQ ID NOS: 1-20) as shown in FIG. 5A. The PPMOs were evaluated for exon skipping efficacy by treating RD cells at various concentrations as described above in the Materials and Methods. Five PPMOs (SEQ ID NOS: 4, 8, 11, 12 and 13) were identified as effective in inducing exon-skipping and selected for additional evaluation. Dose-ranging experiments in RD cells were used to confirm the relative efficacy of these five PPMO sequences as shown in FIGS. 5C to 5H. SEQ ID NOS: 8, 11 and 12 were shown to be most effective at inducing exon 44 skipping as shown in FIG. 5H with SEQ ID NO: 12 proving the most efficacious.

[0295] Comparison of SEQ ID NO: 12 to other exon 44 antisense sequences was done in both RD cells and human primary skeletal muscle cells. All the evaluated sequences were made as peptide-conjugated PPMOs using the CP06062 peptide (SEQ ID NO: 578). This allowed direct comparison of the relative effectiveness of the antisense sequences without regard to antisense chemistry or cell delivery.

[0296] The alignment of the sequences (SEQ ID NOS: 600, 601, 602 and 603) with SEQ ID NOS: 4, 8, 11 and 12 is shown in FIG. 5B. SEQ ID NOS: 601 and 603 are listed as SEQ ID NOS: 165 and 167 in WO2006/000057. SEQ ID NO: 602 is listed in WO2004/083446 and as SEQ ID NO: 21 in U.S. application Ser. No. 11/233,507. SEQ ID NO: 600 was published in 2007 (Wilton, Fall et al. 2007). The comparison in RD cells showed that both SEQ ID NOS: 602 and 603 were superior to SEQ ID NO: 12 (FIG. 5I). However, as shown in FIG. 5J, in human primary skeletal muscle cells SEQ ID NO: 12 was superior (8.86% exon skipping) to SEQ ID NO: 602 (6.42%). Similar experiments are performed with SEQ ID NO: 603.

#### Example 5

##### Exon 45 Scan

[0297] A series of overlapping antisense PPMOs that target human dystrophin exon 45 were designed and synthesized.

For exon 45, a series of 22 PPMOs, each 25 bases in length, were made (SEQ ID NOS: 21, 23, 25, 27, 29, 31, 32, 34, 35, 37, 39, 41, 43 and 45-53) as shown in FIG. 6A. The PPMOs were evaluated for exon skipping efficacy by treating RD cells and human primary skeletal muscle cells at various concentrations as described above in the Materials and Methods. Five PPMOs (SEQ ID NOS: 27, 29, 34, and 39) were identified as effective in inducing exon-skipping and selected for additional evaluation. Dose-ranging experiments in RD cells were used to confirm the relative efficacy of these four PPMO sequences as shown in FIGS. 6C-G and summarized in FIG. 6H. SEQ ID NO: 49 was used as a negative control in these experiments. SEQ ID NOS: 29 and 34 were shown to be most effective at inducing exon 45 skipping as shown in FIG. 6H.

[0298] Comparison of SEQ ID NO: 34 to other exon 45 antisense sequences was done in both RD cells and human primary skeletal muscle cells. All the evaluated sequences were made as peptide-conjugated PPMOs using the CP06062 peptide (SEQ ID NO: 578). This allowed direct comparison of the relative effectiveness of the antisense sequences without regard to antisense chemistry or cell delivery. The alignment of the sequences (SEQ ID NOS: 604, 605, 606 and 607) with SEQ ID NOS: 27, 29, 34 and 39 is shown in FIG. 6B. SEQ ID NOS: 604 and 607 are listed as SEQ ID NOS: 211 and 207 in WO2006/000057, respectively. SEQ ID NOS: 605 and 606 are listed in U.S. application Ser. No. 11/233,507 as SEQ ID NOS: 23 and 1, respectively. The comparison in RD cells showed that SEQ ID NO: 34 was superior to all four sequences evaluated as shown in FIG. 6I. Testing of these compounds in different populations of human primary skeletal muscle cells is performed as described above.

#### SEQUENCE ID LISTING

[0299] Sequences are shown using the nucleotide base symbols common for DNA: A, G, C and T. Other antisense chemistries such as 2'-O-methyl use U in place of T. Any of the bases may be substituted with inosine (I) especially in stretches of three or more G residues.

Name	Sequences	SEQ ID NO.
Oligomer Targeting Sequences (5' to 3'):		
Hu.DMD.Exon44.25.001	CTGCAGGTAAAAGCATATGGATCAA	1
Hu.DMD.Exon44.25.002	ATCGCCTGCAGGTAAAAGCATATGG	2
Hu.DMD.Exon44.25.003	GTCAAATCGCCTGCAGGTAAAAGCA	3
Hu.DMD.Exon44.25.004	GATCTGTCAAATCGCCTGCAGGTAA	4
Hu.DMD.Exon44.25.005	CAACAGATCTGTCAAATCGCCTGCA	5
Hu.DMD.Exon44.25.006	TTTCTCAACAGATCTGTCAAATCGC	6
Hu.DMD.Exon44.25.007	CCATTTCTCAACAGATCTGTCAAAT	7
Hu.DMD.Exon44.25.008	ATAATGAAAACGCCGCAATTTCTCA	8
Hu.DMD.Exon44.25.009	AAATATCTTTATATCATAATGAAAA	9

US 2010/0130591 A1

May 27, 2010

26

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon44.25.010	TGTTAGCCACTGATTAAATATCTTT	10
Hu.DMD.Exon44.25.011	AAACTGTTTCAGCTTCTGTTAGCCAC	11
Hu.DMD.Exon44.25.012	TTGTGTCTTTCTGAGAACTGTTCA	12
Hu.DMD.Exon44.25.013	CCAATTCTCAGGAATTGTGTCTTT	13
Hu.DMD.Exon44.25.014	GTATTTAGCATGTTCCCAATTCTCA	14
Hu.DMD.Exon44.25.015	CTTAAGATACCATTGTATTTAGCA	15
Hu.DMD.Exon44.25.016	CTTACCTTAAGATACCATTGTATT	16
Hu.DMD.Exon44.25.017	AAAGACTTACCTTAAGATACCATT	17
Hu.DMD.Exon44.25.018	AAATCAAAGACTTACCTTAAGATAC	18
Hu.DMD.Exon44.25.019	AAACAAATCAAAGACTTACCTTAA	19
Hu.DMD.Exon44.25.020	TCGAAAAACAAATCAAAGACTTAC	20
Hu.DMD.Exon45.25.001	CTGTAAGATACCAAAAGGCAAAAC	21
Hu.DMD.Exon45.25.002	CCTGTAAGATACCAAAAGGCAAAA	22
Hu.DMD.Exon45.25.002.2	AGTTCCTGTAAGATACCAAAAGGC	23
Hu.DMD.Exon45.25.003	GAGTTCCTGTAAGATACCAAAAGG	24
Hu.DMD.Exon45.25.003.2	CCTGGAGTTCCTGTAAGATACCAA	25
Hu.DMD.Exon45.25.004	TCCTGGAGTTCCTGTAAGATACCAA	26
Hu.DMD.Exon45.25.004.2	GCCATCCTGGAGTTCCTGTAAGATA	27
Hu.DMD.Exon45.25.005	TGCCATCCTGGAGTTCCTGTAAGAT	28
Hu.DMD.Exon45.25.005.2	CCAATGCCATCCTGGAGTTCCTGTA	29
Hu.DMD.Exon45.25.006	CCCAATGCCATCCTGGAGTTCCTGT	30
Hu.DMD.Exon45.25.006.2	GCTGCCCAATGCCATCCTGGAGTTC	31
Hu.DMD.Exon45.25.007	CGCTGCCCAATGCCATCCTGGAGTT	32
Hu.DMD.Exon45.25.008	AACAGTTTGCCGCTGCCCAATGCCA	33
Hu.DMD.Exon45.25.008.2	CTGACAACAGTTTGCCGCTGCCCAA	34
Hu.DMD.Exon45.25.009	GTTGCATTCAATGTTCTGACAACAG	35
Hu.DMD.Exon45.25.010	GCTGAATTATTTCTTCCCAGTTGC	36
Hu.DMD.Exon45.25.010.2	ATTATTTCTTCCCAGTTGCATTCA	37
Hu.DMD.Exon45.25.011	GGCATCTGTTTTTGAGGATTGCTGA	38
Hu.DMD.Exon45.25.011.2	TTTGAGGATTGCTGAATTATTTCTT	39
Hu.DMD.Exon45.25.012	AATTTTCTCTGTAATACTGGCAT	40
Hu.DMD.Exon45.25.012.2	ATACTGGCATCTGTTTTTGAGGATT	41
Hu.DMD.Exon45.25.013	ACCGCAGATTCAGGCTTCCCAATTT	42
Hu.DMD.Exon45.25.013.2	AATTTTCTCTGTAATACTGGCAT	43
Hu.DMD.Exon45.25.014	CTGTTTGACAGCTCCTGCCACCGC	44
Hu.DMD.Exon45.25.014.2	AGATTCAGGCTTCCCAATTTTCTT	45
Hu.DMD.Exon45.25.015	CTCTTTTTTCTGTCTGACAGCTGT	46

US 2010/0130591 A1

May 27, 2010

27

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon45.25.015.2	ACCTCCTGCCACCGCAGATTCAGGC	47
Hu.DMD.Exon45.25.016	CCTACCTCTTTTTTCTGTCTGACAG	48
Hu.DMD.Exon45.25.016.2	GACAGCTGTTTGACAGACCTCCTGCC	49
Hu.DMD.Exon45.25.017	GTCGCCCTACCTCTTTTTTCTGTCT	50
Hu.DMD.Exon45.25.018	GATCTGTGCGCCCTACCTCTTTTTTC	51
Hu.DMD.Exon45.25.019	TATTAGATCTGTGCGCCCTACCTCTT	52
Hu.DMD.Exon45.25.020	ATTCCTATTAGATCTGTGCGCCCTAC	53
Hu.DMD.Exon45.20.001	AGATACCAAAAAGGCAAAAC	54
Hu.DMD.Exon45.20.002	AAGATACCAAAAAGGCAAAA	55
Hu.DMD.Exon45.20.003	CCTGTAAGATACCAAAAAGG	56
Hu.DMD.Exon45.20.004	GAGTTCCTGTAAGATACCAA	57
Hu.DMD.Exon45.20.005	TCCTGGAGTTCCTGTAAGAT	58
Hu.DMD.Exon45.20.006	TGCCATCCTGGAGTTCCTGT	59
Hu.DMD.Exon45.20.007	CCCAATGCCATCCTGGAGTT	60
Hu.DMD.Exon45.20.008	CGCTGCCCAATGCCATCCTG	61
Hu.DMD.Exon45.20.009	CTGACAACAGTTTGCCGCTG	62
Hu.DMD.Exon45.20.010	GTTGCATTCAATGTTCTGAC	63
Hu.DMD.Exon45.20.011	ATTATTTCTTCCCAGTTGC	64
Hu.DMD.Exon45.20.012	TTTGAGGATTGCTGAATTAT	65
Hu.DMD.Exon45.20.013	ATACTGGCATCTGTTTTTGA	66
Hu.DMD.Exon45.20.014	AATTTTCTCTGTAGAATACT	67
Hu.DMD.Exon45.20.015	AGATTTCAGGCTTCCCAATTT	68
Hu.DMD.Exon45.20.016	ACCTCCTGCCACCGCAGATT	69
Hu.DMD.Exon45.20.017	GACAGCTGTTTGACAGACCTC	70
Hu.DMD.Exon45.20.018	CTCTTTTTTCTGTCTGACAG	71
Hu.DMD.Exon45.20.019	CCTACCTCTTTTTTCTGTCT	72
Hu.DMD.Exon45.20.020	GTCGCCCTACCTCTTTTTTC	73
Hu.DMD.Exon45.20.021	GATCTGTGCGCCCTACCTCTT	74
Hu.DMD.Exon45.20.022	TATTAGATCTGTGCGCCCTAC	75
Hu.DMD.Exon45.20.023	ATTCCTATTAGATCTGTCTGC	76
Hu.DMD.Exon46.25.001	GGGGGATTGAGAAAATAAAATTAC	77
Hu.DMD.Exon46.25.002	ATTTGAGAAAATAAAATTACCTTGA	78
Hu.DMD.Exon46.25.002.2	CTAGCCTGGAGAAAGAATAAAA	79
Hu.DMD.Exon46.25.003	AGAAAATAAAATTACCTTGACTTGC	80
Hu.DMD.Exon46.25.003.2	TTCTTCTAGCCTGGAGAAAGAAGAA	81
Hu.DMD.Exon46.25.004	ATAAAATTACCTTGACTTGCTCAAG	82
Hu.DMD.Exon46.25.004.2	TTTTGTTCTTCTAGCCTGGAGAAAG	83

US 2010/0130591 A1

May 27, 2010

28

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon46.25.005	ATTACCTTGACTTGCTCAAGCTTTT	84
Hu.DMD.Exon46.25.005.2	TATTCTTTTGTTCCTCTAGCCTGGA	85
Hu.DMD.Exon46.25.006	CTTGACTTGCTCAAGCTTTTCTTTT	86
Hu.DMD.Exon46.25.006.2	CAAGATATTCTTTTGTTCCTCTAGC	87
Hu.DMD.Exon46.25.007	CTTTTAGTTGCTGCTCTTTTCCAGG	88
Hu.DMD.Exon46.25.008	CCAGGTTCAAGTGGGATACTAGCAA	89
Hu.DMD.Exon46.25.008.2	ATCTCTTTGAAATTCTGACAAGATA	90
Hu.DMD.Exon46.25.009	AGCAATGTTATCTGCTTCCTCCAAC	91
Hu.DMD.Exon46.25.009.2	AACAAATTCATTAAATCTCTTTGA	92
Hu.DMD.Exon46.25.010	CCAACCATAAAACAAATTCATTAA	93
Hu.DMD.Exon46.25.010.2	TTCCTCCAACCATAAAACAAATTC	94
Hu.DMD.Exon46.25.011	TTTAAATCTCTTTGAAATCTGACA	95
Hu.DMD.Exon46.25.012	TGACAAGATATTCTTTTGTTCCTCT	96
Hu.DMD.Exon46.25.012.2	TTCAAGTGGGATACTAGCAATGTTA	97
Hu.DMD.Exon46.25.013	AGATATTCTTTTGTTCCTCTAGCCT	98
Hu.DMD.Exon46.25.013.2	CTGCTCTTTTCCAGGTTCAAGTGGG	99
Hu.DMD.Exon46.25.014	TTCTTTTGTTCCTCTAGCCTGGAGA	100
Hu.DMD.Exon46.25.014.2	CTTTTCTTTTAGTTGCTGCTCTTTT	101
Hu.DMD.Exon46.25.015	TGTTCCTCTAGCCTGGAGAAAGAA	102
Hu.DMD.Exon46.25.016	CTTCTAGCCTGGAGAAAGAAGAATA	103
Hu.DMD.Exon46.25.017	AGCCTGGAGAAAGAAGAATAAAATT	104
Hu.DMD.Exon46.25.018	CTGGAGAAAGAAGAATAAAATTGTT	105
Hu.DMD.Exon46.20.001	GAAAGAAGAATAAAATTGTT	106
Hu.DMD.Exon46.20.002	GGAGAAAGAAGAATAAAATT	107
Hu.DMD.Exon46.20.003	AGCCTGGAGAAAGAAGAATA	108
Hu.DMD.Exon46.20.004	CTTCTAGCCTGGAGAAAGAA	109
Hu.DMD.Exon46.20.005	TTGTTCTTCTAGCCTGGAGA	110
Hu.DMD.Exon46.20.006	TTCTTTTGTTCCTCTAGCCT	111
Hu.DMD.Exon46.20.007	TGACAAGATATTCTTTTGT	112
Hu.DMD.Exon46.20.008	ATCTCTTTGAAATCTGACA	113
Hu.DMD.Exon46.20.009	AACAAATTCATTAAATCTC	114
Hu.DMD.Exon46.20.010	TTCCTCCAACCATAAAACAA	115
Hu.DMD.Exon46.20.011	AGCAATGTTATCTGCTTCCT	116
Hu.DMD.Exon46.20.012	TTCAAGTGGGATACTAGCAA	117
Hu.DMD.Exon46.20.013	CTGCTCTTTTCCAGGTTCAA	118
Hu.DMD.Exon46.20.014	CTTTTCTTTTAGTTGCTGCT	119
Hu.DMD.Exon46.20.015	CTTGACTTGCTCAAGCTTTT	120

US 2010/0130591 A1

May 27, 2010

29

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon46.20.016	ATTACCTTGACTTGCTCAAG	121
Hu.DMD.Exon46.20.017	ATAAAATTACCTTGACTTGC	122
Hu.DMD.Exon46.20.018	AGAAAAATAAAATTACCTTGA	123
Hu.DMD.Exon46.20.019	ATTTGAGAAAATAAAATTAC	124
Hu.DMD.Exon46.20.020	GGGGGATTTGAGAAAATAAA	125
Hu.DMD.Exon47.25.001	CTGAAACAGACAAATGCAACAACGT	126
Hu.DMD.Exon47.25.002	AGTAACTGAAACAGACAAATGCAAC	127
Hu.DMD.Exon47.25.003	CCACCAGTAACTGAAACAGACAAAT	128
Hu.DMD.Exon47.25.004	CTCTTCCACCAGTAACTGAAACAGA	129
Hu.DMD.Exon47.25.005	GGCAACTCTTCCACCAGTAACTGAA	130
Hu.DMD.Exon47.25.006	GCAGGGGCAACTCTTCCACCAGTAA	131
Hu.DMD.Exon47.25.007	CTGGCGCAGGGGCAACTCTTCCACC	132
Hu.DMD.Exon47.25.008	TTTAATTGTTTGAGAATTCCTGGC	133
Hu.DMD.Exon47.25.008.2	TTGTTTGAGAATTCCTGGCGCAGG	134
Hu.DMD.Exon47.25.009	GCACGGGTCCTCCAGTTTCATTTAA	135
Hu.DMD.Exon47.25.009.2	TCCAGTTTCATTTAATTGTTTGAGA	136
Hu.DMD.Exon47.25.010	GCTTATGGGAGCACTTACAAGCACG	137
Hu.DMD.Exon47.25.010.2	TACAAGCACGGGTCCTCCAGTTTCA	138
Hu.DMD.Exon47.25.011	AGTTTATCTTGCTCTTCTGGGCTTA	139
Hu.DMD.Exon47.25.012	TCTGCTTGAGCTTATTTTCAAGTTT	140
Hu.DMD.Exon47.25.012.2	ATCTTGCTCTTCTGGGCTTATGGGA	141
Hu.DMD.Exon47.25.013	CTTTATCCACTGGAGATTGTCTGTC	142
Hu.DMD.Exon47.25.013.2	CTTATTTTCAAGTTTATCTTGCTCT	143
Hu.DMD.Exon47.25.014	CTAACCTTTATCCACTGGAGATTG	144
Hu.DMD.Exon47.25.014.2	ATTTGTCTGCTTGAGCTTATTTTCA	145
Hu.DMD.Exon47.25.015	AATGTCTAACCTTTATCCACTGGAG	146
Hu.DMD.Exon47.25.016	TGGTTAATGTCTAACCTTTATCCAC	147
Hu.DMD.Exon47.25.017	AGAGATGGTTAATGTCTAACCTTTA	148
Hu.DMD.Exon47.25.018	ACGGAAGAGATGGTTAATGTCTAAC	149
Hu.DMD.Exon47.20.001	ACAGACAAATGCAACAACGT	150
Hu.DMD.Exon47.20.002	CTGAAACAGACAAATGCAAC	151
Hu.DMD.Exon47.20.003	AGTAACTGAAACAGACAAAT	152
Hu.DMD.Exon47.20.004	CCACCAGTAACTGAAACAGA	153
Hu.DMD.Exon47.20.005	CTCTTCCACCAGTAACTGAA	154
Hu.DMD.Exon47.20.006	GGCAACTCTTCCACCAGTAA	155
Hu.DMD.Exon47.20.007	CTGGCGCAGGGGCAACTCTT	156
Hu.DMD.Exon47.20.008	TTGTTTGAGAATTCCTGGC	157

US 2010/0130591 A1

May 27, 2010

30

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon47.20.009	TCCAGTTTCATTTAATTGTT	158
Hu.DMD.Exon47.20.010	TACAAGCACGGTCCTCCAG	159
Hu.DMD.Exon47.20.011	GCTTATGGGAGCACTTACAA	160
Hu.DMD.Exon47.20.012	ATCTTGCTCTTCTGGGCTTA	161
Hu.DMD.Exon47.20.013	CTTATTTTCAAGTTTATCTT	162
Hu.DMD.Exon47.20.014	ATTTGTCTGCTTGAGCTTAT	163
Hu.DMD.Exon47.20.015	CTTTATCCACTGGAGATTG	164
Hu.DMD.Exon47.20.016	CTAACCTTTATCCACTGGAG	165
Hu.DMD.Exon47.20.017	AATGTCTAACCTTTATCCAC	166
Hu.DMD.Exon47.20.018	TGGTTAATGTCTAACCTTTA	167
Hu.DMD.Exon47.20.019	AGAGATGGTTAATGTCTAAC	168
Hu.DMD.Exon47.20.020	ACGGAAGAGATGGTTAATGT	169
Hu.DMD.Exon48.25.001	CTGAAAGGAAAATACATTTTAAAA	170
Hu.DMD.Exon48.25.002	CCTGAAAGGAAAATACATTTTAAAA	171
Hu.DMD.Exon48.25.002.2	GAAACCTGAAAGGAAAATACATTTT	172
Hu.DMD.Exon48.25.003	GGAAACCTGAAAGGAAAATACATTT	173
Hu.DMD.Exon48.25.003.2	CTCTGGAACCTGAAAGGAAAATAC	174
Hu.DMD.Exon48.25.004	GCTCTGGAACCTGAAAGGAAAATA	175
Hu.DMD.Exon48.25.004.2	TAAAGCTCTGGAACCTGAAAGGAA	634
Hu.DMD.Exon48.25.005	GTAAAGCTCTGGAACCTGAAAGGA	176
Hu.DMD.Exon48.25.005.2	TCAGGTAAAGCTCTGGAACCTGAA	177
Hu.DMD.Exon48.25.006	CTCAGGTAAAGCTCTGGAACCTGA	178
Hu.DMD.Exon48.25.006.2	GTTTCTCAGGTAAAGCTCTGGAAC	179
Hu.DMD.Exon48.25.007	TGTTTCTCAGGTAAAGCTCTGGAAC	180
Hu.DMD.Exon48.25.007.2	AATTTCTCCTTGTCTCAGGTAA	181
Hu.DMD.Exon48.25.008	TTTGAGCTTCAATTTCTCCTTGTTT	182
Hu.DMD.Exon48.25.008	TTTTATTTGAGCTTCAATTTCTCCT	183
Hu.DMD.Exon48.25.009	AAGCTGCCCAAGGTCTTTTATTGA	184
Hu.DMD.Exon48.25.010	AGGTCTTCAAGCTTTTTTTCAAGCT	185
Hu.DMD.Exon48.25.010.2	TTCAAGCTTTTTTTCAAGCTGCCCA	186
Hu.DMD.Exon48.25.011	GATGATTTAACTGCTCTTCAAGGTC	187
Hu.DMD.Exon48.25.011.2	CTGCTCTTCAAGGTCTTCAAGCTTT	188
Hu.DMD.Exon48.25.012	AGGAGATAACCACAGCAGCAGATGA	189
Hu.DMD.Exon48.25.012.2	CAGCAGATGATTTAACTGCTCTTCA	190
Hu.DMD.Exon48.25.013	ATTTCCAAGTATTCCTAATAGGAG	191
Hu.DMD.Exon48.25.014	CTTGGTTTGGTTGGTTATAAATTC	192
Hu.DMD.Exon48.25.014.2	CAACTGATTCCTAATAGGAGATAAC	193

US 2010/0130591 A1

May 27, 2010

31

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon48.25.015	CTTAACGTCAAATGGTCCTTCTTGG	194
Hu.DMD.Exon48.25.015.2	TTGGTTATAAATTTCCAACCTGATTC	195
Hu.DMD.Exon48.25.016	CCTACCTTAACGTCAAATGGTCCTT	196
Hu.DMD.Exon48.25.016.2	TCCTTCTTGGTTTGGTTGGTTATAA	197
Hu.DMD.Exon48.25.017	AGTTCCCTACCTTAACGTCAAATGG	198
Hu.DMD.Exon48.25.018	CAAAAAGTTCCCTACCTTAACGTCA	199
Hu.DMD.Exon48.25.019	TAAAGCAAAAAGTTCCCTACCTTAA	200
Hu.DMD.Exon48.25.020	ATATTTAAAGCAAAAAGTTCCCTAC	201
Hu.DMD.Exon48.20.001	AGGAAAATACATTTTAAAAA	202
Hu.DMD.Exon48.20.002	AAGGAAAATACATTTTAAAAA	203
Hu.DMD.Exon48.20.003	CCTGAAAGGAAAATACATTT	204
Hu.DMD.Exon48.20.004	GGAAACCTGAAAGGAAAATA	205
Hu.DMD.Exon48.20.005	GCTCTGGAAACCTGAAAGGA	206
Hu.DMD.Exon48.20.006	GTAAAGCTCTGGAAACCTGA	207
Hu.DMD.Exon48.20.007	CTCAGGTAAAGCTCTGGAAA	208
Hu.DMD.Exon48.20.008	AATTTCTCCTTGTTTCTCAG	209
Hu.DMD.Exon48.20.009	TTTTATTTGAGCTTCAATTT	210
Hu.DMD.Exon48.20.010	AAGCTGCCCAAGGTCTTTTA	211
Hu.DMD.Exon48.20.011	TTCAAGCTTTTTTTCAAGCT	212
Hu.DMD.Exon48.20.012	CTGCTCTTCAAGGTCTTCAA	213
Hu.DMD.Exon48.20.013	CAGCAGATGATTTAACTGCT	214
Hu.DMD.Exon48.20.014	AGGAGATAACCACAGCAGCA	215
Hu.DMD.Exon48.20.015	CAACTGATTCCTAATAGGAG	216
Hu.DMD.Exon48.20.016	TTGGTTATAAATTTCCAACCT	217
Hu.DMD.Exon48.20.017	TCCTTCTTGGTTTGGTTGGT	218
Hu.DMD.Exon48.20.018	CTTAACGTCAAATGGTCCTT	219
Hu.DMD.Exon48.20.019	CCTACCTTAACGTCAAATGG	220
Hu.DMD.Exon48.20.020	AGTTCCCTACCTTAACGTCA	221
Hu.DMD.Exon48.20.021	CAAAAAGTTCCCTACCTTAA	222
Hu.DMD.Exon48.20.022	TAAAGCAAAAAGTTCCCTAC	223
Hu.DMD.Exon48.20.023	ATATTTAAAGCAAAAAGTTC	224
Hu.DMD.Exon49.25.001	CTGGGGAAAAGAACCCATATAGTGC	225
Hu.DMD.Exon49.25.002	TCCTGGGGAAAAGAACCCATATAGT	226
Hu.DMD.Exon49.25.002.2	GTTTCCTGGGGAAAAGAACCCATAT	227
Hu.DMD.Exon49.25.003	CAGTTTCCTGGGGAAAAGAACCCAT	228
Hu.DMD.Exon49.25.003.2	TTTCAGTTTCCTGGGGAAAAGAACC	229
Hu.DMD.Exon49.25.004	TATTTTCAGTTTCCTGGGGAAAAGAA	230



US 2010/0130591 A1

May 27, 2010

32

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon49.25.004.2	TGCTATTTTCAGTTTCCTGGGGAAAA	231
Hu.DMD.Exon49.25.005	ACTGCTATTTTCAGTTTCCTGGGGAA	232
Hu.DMD.Exon49.25.005.2	TGAACTGCTATTTTCAGTTTCCTGGG	233
Hu.DMD.Exon49.25.006	CTTGAAC TGCTATTTTCAGTTTCCTG	234
Hu.DMD.Exon49.25.006.2	TAGCTTGAAC TGCTATTTTCAGTTTC	235
Hu.DMD.Exon49.25.007	TTTAGCTTGAAC TGCTATTTTCAGTT	236
Hu.DMD.Exon49.25.008	TTCCACATCCGTTGTTTAGCTTGA	237
Hu.DMD.Exon49.25.009	TGCCCTTTAGACAAAATCTCTTCCA	238
Hu.DMD.Exon49.25.009.2	TTTAGACAAAATCTCTTCCACATCC	239
Hu.DMD.Exon49.25.010	GTTTTTCCTTGTACAAATGCTGCCC	240
Hu.DMD.Exon49.25.010.2	GTACAAATGCTGCCCTTTAGACAAA	241
Hu.DMD.Exon49.25.011	CTTCACTGGCTGAGTGGCTGGTTTT	242
Hu.DMD.Exon49.25.011.2	GGCTGGTTTTTCCTTGTACAAATGC	243
Hu.DMD.Exon49.25.012	ATTACCTTCACTGGCTGAGTGGCTG	244
Hu.DMD.Exon49.25.013	GCTTCATTACCTTCACTGGCTGAGT	245
Hu.DMD.Exon49.25.014	AGGTTGCTTCATTACCTTCACTGGC	246
Hu.DMD.Exon49.25.015	GCTAGAGGTTGCTTCATTACCTTCA	247
Hu.DMD.Exon49.25.016	ATATTGCTAGAGGTTGCTTCATTAC	248
Hu.DMD.Exon49.20.001	GAAAAGAACCCATATAGTGC	249
Hu.DMD.Exon49.20.002	GGGAAAAGAACCCATATAGT	250
Hu.DMD.Exon49.20.003	TCCTGGGGAAAAGAACCCAT	251
Hu.DMD.Exon49.20.004	CAGTTTCCTGGGGAAAAGAA	252
Hu.DMD.Exon49.20.005	TATTTTCAGTTTCCTGGGGAA	253
Hu.DMD.Exon49.20.006	ACTGCTATTTTCAGTTTCCTG	254
Hu.DMD.Exon49.20.007	CTTGAAC TGCTATTTTCAGTT	255
Hu.DMD.Exon49.20.008	TTTAGCTTGAAC TGCTATTT	256
Hu.DMD.Exon49.20.009	TTCCACATCCGTTGTTTAG	257
Hu.DMD.Exon49.20.010	TTTAGACAAAATCTCTTCCA	258
Hu.DMD.Exon49.20.011	GTACAAATGCTGCCCTTTAG	259
Hu.DMD.Exon49.20.012	GGCTGGTTTTTCCTTGTACA	260
Hu.DMD.Exon49.20.013	CTTCACTGGCTGAGTGGCTG	261
Hu.DMD.Exon49.20.014	ATTACCTTCACTGGCTGAGT	262
Hu.DMD.Exon49.20.015	GCTTCATTACCTTCACTGGC	263
Hu.DMD.Exon49.20.016	AGGTTGCTTCATTACCTTCA	264
Hu.DMD.Exon49.20.017	GCTAGAGGTTGCTTCATTAC	265
Hu.DMD.Exon49.20.018	ATATTGCTAGAGGTTGCTTC	266
Hu.DMD.Exon50.25.001	CTTTAACAGAAAAGCATACATTA	267

US 2010/0130591 A1

May 27, 2010

33

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon50.25.002	TCCTCTTTAACAGAAAAGCATACAC	268
Hu.DMD.Exon50.25.002.2	TTCTCTTTAACAGAAAAGCATACA	269
Hu.DMD.Exon50.25.003	TAACCTCCTCTTTAACAGAAAAGCA	270
Hu.DMD.Exon50.25.003.2	CTAACTTCCTCTTTAACAGAAAAGC	271
Hu.DMD.Exon50.25.004	TCTTCTAACTTCCTCTTTAACAGAA	272
Hu.DMD.Exon50.25.004.2	ATCTTCTAACTTCCTCTTTAACAGA	273
Hu.DMD.Exon50.25.005	TCAGATCTTCTAACTTCCTCTTTAA	274
Hu.DMD.Exon50.25.005.2	CTCAGATCTTCTAACTTCCTCTTTA	275
Hu.DMD.Exon50.25.006	AGAGCTCAGATCTTCTAACTTCCTC	276
Hu.DMD.Exon50.25.006.2	CAGAGCTCAGATCTTCTAACTTCCT	277
NG-08-0731		
Hu.DMD.Exon50.25.007	CACTCAGAGCTCAGATCTTCTACT	278
Hu.DMD.Exon50.25.007.2	CCTTCCACTCAGAGCTCAGATCTTC	279
Hu.DMD.Exon50.25.008	GTAAACGGTTTACCGCCTTCCACTC	280
Hu.DMD.Exon50.25.009	CTTTGCCCTCAGCTCTTGAAGTAAA	281
Hu.DMD.Exon50.25.009.2	CCCTCAGCTCTTGAAGTAAACGGTT	282
Hu.DMD.Exon50.25.010	CCAGGAGCTAGGTCAGGCTGCTTTG	283
Hu.DMD.Exon50.25.010.2	GGTCAGGCTGCTTTGCCCTCAGCTC	284
Hu.DMD.Exon50.25.011	AGGCTCCAATAGTGGTCAGTCCAGG	285
Hu.DMD.Exon50.25.011.2	TCAGTCCAGGAGCTAGGTGAGGCTG	286
Hu.DMD.Exon50.25.012	CTTACAGGCTCCAATAGTGGTCAGT	287
AVI-5038		
Hu.DMD.Exon50.25.013	GTATACTTACAGGCTCCAATAGTGG	288
Hu.DMD.Exon50.25.014	ATCCAGTATACTTACAGGCTCCAAT	289
Hu.DMD.Exon50.25.015	ATGGGATCCAGTATACTTACAGGCT	290
NG-08-0741		
Hu.DMD.Exon50.25.016	AGAGAATGGGATCCAGTATACTTAC	291
NG-08-0742		
Hu.DMD.Exon50.20.001	ACAGAAAAGCATACACATTA	292
Hu.DMD.Exon50.20.002	TTTAACAGAAAAGCATACAC	293
Hu.DMD.Exon50.20.003	TCCTCTTTAACAGAAAAGCA	294
Hu.DMD.Exon50.20.004	TAACCTCCTCTTTAACAGAA	295
Hu.DMD.Exon50.20.005	TCTTCTAACTTCCTCTTTAA	296
Hu.DMD.Exon50.20.006	TCAGATCTTCTAACTTCCTC	297
Hu.DMD.Exon50.20.007	CCTTCCACTCAGAGCTCAGA	298
Hu.DMD.Exon50.20.008	GTAAACGGTTTACCGCCTTC	299
Hu.DMD.Exon50.20.009	CCCTCAGCTCTTGAAGTAAA	300
Hu.DMD.Exon50.20.010	GGTCAGGCTGCTTTGCCCTC	301
Hu.DMD.Exon50.20.011	TCAGTCCAGGAGCTAGGTCA	302

US 2010/0130591 A1

May 27, 2010

34

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon50.20.012	AGGCTCCAATAGTGGTCAGT	303
Hu.DMD.Exon50.20.013	CTTACAGGCTCCAATAGTGG	304
Hu.DMD.Exon50.20.014	GTATACTTACAGGCTCCAAT	305
Hu.DMD.Exon50.20.015	ATCCAGTATACTTACAGGCT	306
Hu.DMD.Exon50.20.016	ATGGGATCCAGTATACTTAC	307
Hu.DMD.Exon50.20.017	AGAGAATGGGATCCAGTATA	308
Hu.DMD.Exon51.25.001-44	CTAAAATATTTTGGGTTTTTGCAAAA	309
Hu.DMD.Exon51.25.002-45	GCTAAAATATTTTGGGTTTTTGCAAA	310
Hu.DMD.Exon51.25.002.2-46	TAGGAGCTAAAATATTTTGGGTTTTT	311
Hu.DMD.Exon51.25.003	AGTAGGAGCTAAAATATTTTGGGTT	312
Hu.DMD.Exon51.25.003.2	TGAGTAGGAGCTAAAATATTTTGGG	313
Hu.DMD.Exon51.25.004	CTGAGTAGGAGCTAAAATATTTTGGG	314
Hu.DMD.Exon51.25.004.2	CAGTCTGAGTAGGAGCTAAAATATT	315
Hu.DMD.Exon51.25.005	ACAGTCTGAGTAGGAGCTAAAATATT	316
Hu.DMD.Exon51.25.005.2	GAGTAACAGTCTGAGTAGGAGCTAAA	317
Hu.DMD.Exon51.25.006	CAGAGTAACAGTCTGAGTAGGAGCT	318
Hu.DMD.Exon51.25.006.2	CACCAGAGTAACAGTCTGAGTAGGAG	319
Hu.DMD.Exon51.25.007	GTCACCAGAGTAACAGTCTGAGTAG	320
Hu.DMD.Exon51.25.007.2	AACCACAGGTTGTGTACCAGAGTAA	321
Hu.DMD.Exon51.25.008	GTTGTGTACCAGAGTAACAGTCTG	322
Hu.DMD.Exon51.25.009	TGGCAGTTTCCTTAGTAACCACAGGT	323
Hu.DMD.Exon51.25.010	ATTTCTAGTTTGGAGATGGCAGTTTC	324
Hu.DMD.Exon51.25.010.2	GGAAGATGGCATTCTAGTTTGGAG	325
Hu.DMD.Exon51.25.011	CATCAAGGAAGATGGCATTCTAGTT	326
Hu.DMD.Exon51.25.011.2	GAGCAGGTACCTCCAACATCAAGGAA	327
Hu.DMD.Exon51.25.012	ATCTGCCAGAGCAGGTACCTCCAAC	328
Hu.DMD.Exon51.25.013	AAGTTCTGTCCAAGCCCGTTGAAAT	329
Hu.DMD.Exon51.25.013.2	CGGTTGAAATCTGCCAGAGCAGGTAC	330
Hu.DMD.Exon51.25.014	GAGAAAGCCAGTCGGTAAGTTCTGTC	331
Hu.DMD.Exon51.25.014.2	GTCGGTAAGTTCTGTCCAAGCCCGG	332
Hu.DMD.Exon51.25.015	ATAACTTGATCAAGCAGAGAAAGCCA	333
Hu.DMD.Exon51.25.015.2	AAGCAGAGAAAGCCAGTCGGTAAGT	334
Hu.DMD.Exon51.25.016	CACCCTCTGTGATTTTATAACTTGAT	335
Hu.DMD.Exon51.25.017	CAAGGTCACCCACCATCACCTCTGT	336
Hu.DMD.Exon51.25.017.2	CATCACCTCTGTGATTTTATAACT	337
Hu.DMD.Exon51.25.018	CTTCTGCTTGATGATCATCTCGTTGA	338

US 2010/0130591 A1

May 27, 2010

35

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon51.25.019	CCTTCTGCTTGATGATCATCTCGTTG	339
Hu.DMD.Exon51.25.019.2	ATCTCGTTGATATCCTCAAGTCACC	340
Hu.DMD.Exon51.25.020	TCATACCTTCTGCTTGATGATCATCT	341
Hu.DMD.Exon51.25.020.2	TCATTTTTTCTCATACCTTCTGCTTG	342
Hu.DMD.Exon51.25.021	TTTTCTCATACCTTCTGCTTGATGAT	343
Hu.DMD.Exon51.25.022	TTTTATCATTTTTTCTCATACCTTCT	344
Hu.DMD.Exon51.25.023	CCAACTTTTATCATTTTTTCTCATAC	345
Hu.DMD.Exon51.20.001	ATATTTTGGGTTTTTGCAAA	346
Hu.DMD.Exon51.20.002	AAAATATTTTGGGTTTTTGC	347
Hu.DMD.Exon51.20.003	GAGCTAAAATATTTTGGGTT	348
Hu.DMD.Exon51.20.004	AGTAGGAGCTAAAATATTTT	349
Hu.DMD.Exon51.20.005	GTCTGAGTAGGAGCTAAAAT	350
Hu.DMD.Exon51.20.006	TAACAGTCTGAGTAGGAGCT	351
Hu.DMD.Exon51.20.007	CAGAGTAACAGTCTGAGTAG	352
Hu.DMD.Exon51.20.008	CACAGGTGTGTCAACAGAG	353
Hu.DMD.Exon51.20.009	AGTTTCCTTAGTAACACAG	354
Hu.DMD.Exon51.20.010	TAGTTTGGAGATGGCAGTTT	355
Hu.DMD.Exon51.20.011	GGAAGATGGCATTCTAGTT	356
Hu.DMD.Exon51.20.012	TACCTCCAACATCAAGGAAG	357
Hu.DMD.Exon51.20.013	ATCTGCCAGAGCAGGTACCT	358
Hu.DMD.Exon51.20.014	CCAAGCCCGGTTGAAATCTG	359
Hu.DMD.Exon51.20.015	GTCGGTAAGTTCTGTCCAAG	360
Hu.DMD.Exon51.20.016	AAGCAGAGAAAGCCAGTCGG	361
Hu.DMD.Exon51.20.017	TTTTATACTTGATCAAGCA	362
Hu.DMD.Exon51.20.018	CATCACCTCTGTGATTTTA	363
Hu.DMD.Exon51.20.019	CTCAAGGTCACCCACCATCA	364
Hu.DMD.Exon51.20.020	CATCTCGTTGATATCCTCAA	365
Hu.DMD.Exon51.20.021	CTTCTGCTTGATGATCATCT	366
Hu.DMD.Exon51.20.022	CATACCTTCTGCTTGATGAT	367
Hu.DMD.Exon51.20.023	TTTCTCATACCTTCTGCTTG	368
Hu.DMD.Exon51.20.024	CATTTTTTCTCATACCTTCT	369
Hu.DMD.Exon51.20.025	TTTATCATTTTTTCTCATAC	370
Hu.DMD.Exon51.20.026	CAACTTTTATCATTTTTTCT	371
Hu.DMD.Exon52.25.001	CTGTAAGAACAAATATCCCTTAGTA	372
Hu.DMD.Exon52.25.002	TGCCTGTAAGAACAAATATCCCTTA	373
Hu.DMD.Exon52.25.002.2	GTTGCCTGTAAGAACAAATATCCCT	374
Hu.DMD.Exon52.25.003	ATTGTTGCCTGTAAGAACAAATATC	375

US 2010/0130591 A1

May 27, 2010

36

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon52.25.003.2	GCATTGTTGCCTGTAAGAACAAATA	376
Hu.DMD.Exon52.25.004	CCTGCATTGTTGCCTGTAAGAACAA	377
Hu.DMD.Exon52.25.004.2	ATCCTGCATTGTTGCCTGTAAGAAC	378
Hu.DMD.Exon52.25.005	CAAATCCTGCATTGTTGCCTGTAAG	379
Hu.DMD.Exon52.25.005.2	TCCAAATCCTGCATTGTTGCCTGTA	380
Hu.DMD.Exon52.25.006	TGTTCCAAATCCTGCATTGTTGCCT	381
Hu.DMD.Exon52.25.006.2	TCTGTTCCAAATCCTGCATTGTTGC	382
Hu.DMD.Exon52.25.007	AACTGGGACGCCTCTGTTCCAAAT	383
Hu.DMD.Exon52.25.007.2	GCCTCTGTTCCAAATCCTGCATTGT	384
Hu.DMD.Exon52.25.008	CAGCGGTAATGAGTTCTTCCAACTG	385
Hu.DMD.Exon52.25.008.2	CTTCCAACTGGGACGCCTCTGTTC	386
Hu.DMD.Exon52.25.009	CTTGTTTTTCAAATTTGGGCAGCG	387
Hu.DMD.Exon52.25.010	CTAGCCTCTTGATTGCTGGTCTTGT	388
Hu.DMD.Exon52.25.010.2	TTTTCAAATTTTGGGCAGCGTAAT	389
Hu.DMD.Exon52.25.011	TTCGATCCGTAATGATTGTTCTAGC	390
Hu.DMD.Exon52.25.011.2	GATTGCTGGTCTTGTTTTTCAAATT	391
Hu.DMD.Exon52.25.012	CTTACTTCGATCCGTAATGATTGTT	392
Hu.DMD.Exon52.25.012.2	TTGTTCTAGCCTCTTGATTGCTGGT	393
Hu.DMD.Exon52.25.013	AAAACTTACTTCGATCCGTAATGA	394
Hu.DMD.Exon52.25.014	TGTTAAAAAATACTTCGATCCGT	395
Hu.DMD.Exon52.25.015	ATGCTTGTTAAAAAATACTTCGA	396
Hu.DMD.Exon52.25.016	GTCCCATGCTTGTTAAAAAATACTAC	397
Hu.DMD.Exon52.20.001	AGAACAATATCCCTTAGTA	398
Hu.DMD.Exon52.20.002	GTAAGAACAATATCCCTTA	399
Hu.DMD.Exon52.20.003	TGCCTGTAAGAACAATATC	400
Hu.DMD.Exon52.20.004	ATTGTTGCCTGTAAGAACAA	401
Hu.DMD.Exon52.20.005	CCTGCATTGTTGCCTGTAAG	402
Hu.DMD.Exon52.20.006	CAAATCCTGCATTGTTGCCT	403
Hu.DMD.Exon52.20.007	GCCTCTGTTCCAAATCCTGC	404
Hu.DMD.Exon52.20.008	CTTCCAACTGGGACGCCTC	405
Hu.DMD.Exon52.20.009	CAGCGGTAATGAGTTCTTCC	406
Hu.DMD.Exon52.20.010	TTTTCAAATTTTGGGCAGCG	407
Hu.DMD.Exon52.20.011	GATTGCTGGTCTTGTTTTTC	408
Hu.DMD.Exon52.20.012	TTGTTCTAGCCTCTTGATTG	409
Hu.DMD.Exon52.20.013	TTCGATCCGTAATGATTGTT	410
Hu.DMD.Exon52.20.014	CTTACTTCGATCCGTAATGA	411
Hu.DMD.Exon52.20.015	AAAACTTACTTCGATCCGT	412

US 2010/0130591 A1

May 27, 2010

37

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon52.20.016	TGTTAAAAAACTTACTTCGA	413
Hu.DMD.Exon52.20.017	ATGCTTGTTAAAAAATTAC	414
Hu.DMD.Exon52.20.018	GTCCCATGCTTGTTAAAAAA	415
Hu.DMD.Exon53.25.001	CTAGAATAAAAGGAAAAATAATAT	416
Hu.DMD.Exon53.25.002	AACTAGAATAAAAGGAAAAATAAT	417
Hu.DMD.Exon53.25.002.2	TTCAACTAGAATAAAAGGAAAAATA	418
Hu.DMD.Exon53.25.003	CTTTCAACTAGAATAAAAGGAAAA	419
Hu.DMD.Exon53.25.003.2	ATTCTTTCAACTAGAATAAAAGGAA	420
Hu.DMD.Exon53.25.004	GAATTCTTTCAACTAGAATAAAAGG	421
Hu.DMD.Exon53.25.004.2	TCTGAATTCTTTCAACTAGAATAAA	422
Hu.DMD.Exon53.25.005	ATTCTGAATTCTTTCAACTAGAATA	423
Hu.DMD.Exon53.25.005.2	CTGATTCTGAATTCTTTCAACTAGA	424
Hu.DMD.Exon53.25.006	CACTGATTCTGAATTCTTTCAACTA	425
Hu.DMD.Exon53.25.006.2	TCCCACTGATTCTGAATTCTTTCAA	426
Hu.DMD.Exon53.25.007	CATCCCACTGATTCTGAATTCTTTTC	427
Hu.DMD.Exon53.25.008	TACTTCATCCCACTGATTCTGAATT	428
Hu.DMD.Exon53.25.008.2	CTGAAGGTGTTCTTGTACTTCATCC	429
Hu.DMD.Exon53.25.009	CGGTTCTGAAGGTGTTCTTGTACT	430
Hu.DMD.Exon53.25.009.2	CTGTTGCCTCCGGTTCTGAAGGTGT	431
Hu.DMD.Exon53.25.010	TTTCATTCAACTGTTGCCTCCGGTT	432
Hu.DMD.Exon53.25.010.2	TAACATTTCAATCAACTGTTGCCTC	433
Hu.DMD.Exon53.25.011	TTGTGTTGAATCCTTTAACATTTC	434
Hu.DMD.Exon53.25.012	TCTTCCTTAGCTTCCAGCCATTGTG	435
Hu.DMD.Exon53.25.012.2	CTTAGCTTCCAGCCATTGTGTTGAA	436
Hu.DMD.Exon53.25.013	GTCCTAAGACCTGCTCAGCTTCTTC	437
Hu.DMD.Exon53.25.013.2	CTGCTCAGCTTCTTCCTTAGCTTCC	438
Hu.DMD.Exon53.25.014	CTCAAGCTTGGCTCTGGCCTGTCCCT	439
Hu.DMD.Exon53.25.014.2	GGCCTGTCTAAGACCTGCTCAGCT	440
Hu.DMD.Exon53.25.015	TAGGGACCTCCTTCCATGACTCAA	441
Hu.DMD.Exon53.25.016	TTTGGATTGCATCTACTGTATAGGG	442
Hu.DMD.Exon53.25.016.2	ACCTCCTTCCATGACTCAAGCTTG	443
Hu.DMD.Exon53.25.017	CTTGGTTTCTGTGATTTTCTTTTG	444
Hu.DMD.Exon53.25.017.2	ATCTACTGTATAGGGACCTCCTTC	445
Hu.DMD.Exon53.25.018	CTAACCTTGGTTTCTGTGATTTTCT	446
Hu.DMD.Exon53.25.018.2	TTTCTTTTGGATTGCATCTACTGTA	447
Hu.DMD.Exon53.25.019	TGATACTAACCTTGGTTTCTGTGAT	448
Hu.DMD.Exon53.25.020	ATCTTTGATACTAACCTTGGTTTCT	449

US 2010/0130591 A1

May 27, 2010

38

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon53.25.021	AAGGTATCTTTGATACTAACCTTGG	450
Hu.DMD.Exon53.25.022	TTAAAAAGGTATCTTTGATACTAAC	451
Hu.DMD.Exon53.20.001	ATAAAAGGAAAAATAAATAT	452
Hu.DMD.Exon53.20.002	GAATAAAAGGAAAAATAAAT	453
Hu.DMD.Exon53.20.003	AACTAGAATAAAAGGAAAA	454
Hu.DMD.Exon53.20.004	CTTTCACCTAGAATAAAAGG	455
Hu.DMD.Exon53.20.005	GAATTCTTTCACCTAGAATA	456
Hu.DMD.Exon53.20.006	ATTCTGAATTCTTTCACCTA	457
Hu.DMD.Exon53.20.007	TACTTCATCCCACTGATTCT	458
Hu.DMD.Exon53.20.008	CTGAAGGTGTTCTTGTA	459
Hu.DMD.Exon53.20.009	CTGTTGCCTCCGGTTCTGAA	460
Hu.DMD.Exon53.20.010	TAACATTTCACTCACTGTT	461
Hu.DMD.Exon53.20.011	TTGTGTTGAATCCTTTAACA	462
Hu.DMD.Exon53.20.012	CTTAGCTTCCAGCCATTGTG	463
Hu.DMD.Exon53.20.013	CTGCTCAGCTTCTTCCTTAG	464
Hu.DMD.Exon53.20.014	GGCCTGTCTTAAGACCTGCT	465
Hu.DMD.Exon53.20.015	CTCAAGCTTGGCTCTGGCCT	466
Hu.DMD.Exon53.20.016	ACCCTCCTTCCATGACTCAA	467
Hu.DMD.Exon53.20.017	ATCTACTGTATAGGACCT	468
Hu.DMD.Exon53.20.018	TTCTTTTGGATTGCATCTA	469
Hu.DMD.Exon53.20.019	CTTGTTTCTGTGATTTTCT	470
Hu.DMD.Exon53.20.020	CTAACCTTGGTTTCTGTGAT	471
Hu.DMD.Exon53.20.021	TGATACTAACCTTGGTTTCT	472
Hu.DMD.Exon53.20.022	ATCTTTGATACTAACCTTGG	473
Hu.DMD.Exon53.20.023	AAGGTATCTTTGATACTAAC	474
Hu.DMD.Exon53.20.024	TTAAAAAGGTATCTTTGATA	475
Hu.DMD.Exon54.25.001	CTATAGATTTTATGAGAAAGAGA	476
Hu.DMD.Exon54.25.002	AACTGCTATAGATTTTATGAGAAA	477
Hu.DMD.Exon54.25.003	TGGCCAACTGCTATAGATTTTATG	478
Hu.DMD.Exon54.25.004	GTCTTTGGCCAACTGCTATAGATTT	479
Hu.DMD.Exon54.25.005	CGGAGGTCTTTGGCCAACTGCTATA	480
Hu.DMD.Exon54.25.006	ACTGGCGGAGGTCTTTGGCCAACTG	481
Hu.DMD.Exon54.25.007	TTTGTCTGCCACTGGCGGAGGTCTT	482
Hu.DMD.Exon54.25.008	AGTCATTTGCCACATCTACATTTGT	483
Hu.DMD.Exon54.25.008.2	TTTGCCACATCTACATTTGTCTGCC	484
Hu.DMD.Exon54.25.009	CCGGAGAAGTTTCAGGGCCAAGTCA	485
Hu.DMD.Exon54.25.010	GTATCATCTGCAGAATAATCCCGGA	486



US 2010/0130591 A1

May 27, 2010

39

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon54.25.010.2	TAATCCCGGAGAAGTTTCAGGGCCA	487
Hu.DMD.Exon54.25.011	TTATCATGTGGACTTTTCTGGTATC	488
Hu.DMD.Exon54.25.012	AGAGGCATTGATATTCTCTGTTATC	489
Hu.DMD.Exon54.25.012.2	ATGTGGACTTTTCTGGTATCATCTG	490
Hu.DMD.Exon54.25.013	CTTTTATGAATGCTTCTCCAAGAGG	491
Hu.DMD.Exon54.25.013.2	ATATTCTCTGTTATCATGTGGACTT	492
Hu.DMD.Exon54.25.014	CATACCTTTTATGAATGCTTCTCCA	493
Hu.DMD.Exon54.25.014.2	CTCCAAGAGGCATTGATATTCTCTG	494
Hu.DMD.Exon54.25.015	TAATTCATACCTTTTATGAATGCTT	495
Hu.DMD.Exon54.25.015.2	CTTTTATGAATGCTTCTCCAAGAGG	496
Hu.DMD.Exon54.25.016	TAATGTAATTCATACCTTTTATGAA	497
Hu.DMD.Exon54.25.017	AGAAATAATGTAATTCATACCTTTT	498
Hu.DMD.Exon54.25.018	GTTTTAGAAATAATGTAATTCATAC	499
Hu.DMD.Exon54.20.001	GATTTTATGAGAAAGAGA	500
Hu.DMD.Exon54.20.002	CTATAGATTTTATGAGAAA	501
Hu.DMD.Exon54.20.003	AACGCTATAGATTTTATG	502
Hu.DMD.Exon54.20.004	TGGCCAACGCTATAGATTT	503
Hu.DMD.Exon54.20.005	GTCTTGGCCAACGCTATA	504
Hu.DMD.Exon54.20.006	CGGAGGCTTTGGCCAACG	505
Hu.DMD.Exon54.20.007	TTGTCTGCCACTGGCGGAG	506
Hu.DMD.Exon54.20.008	TTTGCCACATCTACATTTGT	507
Hu.DMD.Exon54.20.009	TTCAGGGCCAAGTCATTTGC	508
Hu.DMD.Exon54.20.010	TAATCCCGGAGAAGTTTCAG	509
Hu.DMD.Exon54.20.011	GTATCATCTGCAGAATAATC	510
Hu.DMD.Exon54.20.012	ATGTGGACTTTTCTGGTATC	511
Hu.DMD.Exon54.20.013	ATATTCTCTGTTATCATGTG	512
Hu.DMD.Exon54.20.014	CTCCAAGAGGCATTGATATT	513
Hu.DMD.Exon54.20.015	CTTTTATGAATGCTTCTCCA	514
Hu.DMD.Exon54.20.016	CATACCTTTTATGAATGCTT	515
Hu.DMD.Exon54.20.017	TAATTCATACCTTTTATGAA	516
Hu.DMD.Exon54.20.018	TAATGTAATTCATACCTTTT	517
Hu.DMD.Exon54.20.019	AGAAATAATGTAATTCATAC	518
Hu.DMD.Exon54.20.020	GTTTTAGAAATAATGTAATT	519
Hu.DMD.Exon55.25.001	CTGCAAAGGACCAAATGTTTCAGATG	520
Hu.DMD.Exon55.25.002	TCACCCTGCAAAGGACCAAATGTTT	521
Hu.DMD.Exon55.25.003	CTCACTCACCTGCAAAGGACCAAA	522
Hu.DMD.Exon55.25.004	TCTCGCTCACTCACCTGCAAAGGA	523

US 2010/0130591 A1

May 27, 2010

40

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon55.25.005	CAGCCTCTCGCTCACTCACCTGCA	524
Hu.DMD.Exon55.25.006	CAAAGCAGCCTCTCGCTCACTCACC	525
Hu.DMD.Exon55.25.007	TCTTCCAAGCAGCCTCTCGCTCAC	526
Hu.DMD.Exon55.25.007.2	TCTATGAGTTTCTTCCAAGCAGCC	527
Hu.DMD.Exon55.25.008	GTTGCAGTAATCTATGAGTTTCTTC	528
Hu.DMD.Exon55.25.008.2	GAACTGTTGCAGTAATCTATGAGTT	529
Hu.DMD.Exon55.25.009	TTCCAGGTCCAGGGGAACTGTTGC	530
Hu.DMD.Exon55.25.010	GTAAGCCAGGCAAGAACTTTTCCA	531
Hu.DMD.Exon55.25.010.2	CCAGGCAAGAACTTTTCCAGGTCC	532
Hu.DMD.Exon55.25.011	TGGCAGTTGTTTCAGCTTCTGTAAG	533
Hu.DMD.Exon55.25.011.2	TTCAGCTTCTGTAAGCCAGGCAAGA	635
Hu.DMD.Exon55.25.012	GGTAGCATCCTGTAGGACATTGGCA	534
Hu.DMD.Exon55.25.012.2	GACATTGGCAGTTGTTTCAGCTTCT	535
Hu.DMD.Exon55.25.013	TCTAGGAGCCTTTCCTTACGGGTAG	536
Hu.DMD.Exon55.25.014	CTTTTACTCCCTTGGAGTCTTCTAG	537
Hu.DMD.Exon55.25.014.2	GAGCCTTTCCTTACGGGTAGCATCC	538
Hu.DMD.Exon55.25.015	TTGCCATTGTTTCATCAGCTCTTTT	539
Hu.DMD.Exon55.25.015.2	CTTGAGTCTTCTAGGAGCCTTTCC	540
Hu.DMD.Exon55.25.016	CTTACTTGCCATTGTTTCATCAGCT	541
Hu.DMD.Exon55.25.016.2	CAGCTCTTTTACTCCCTTGGAGTCT	542
Hu.DMD.Exon55.25.017	CCTGACTTACTTGCCATTGTTTCAT	543
Hu.DMD.Exon55.25.018	AAATGCCTGACTTACTTGCCATTGT	544
Hu.DMD.Exon55.25.019	AGCGGAAATGCCTGACTTACTTGCC	545
Hu.DMD.Exon55.25.020	GCTAAAGCGGAAATGCCTGACTTAC	546
Hu.DMD.Exon55.20.001	AAGGACCAAATGTTTCAGATG	547
Hu.DMD.Exon55.20.002	CTGCAAAGGACCAAATGTTTC	548
Hu.DMD.Exon55.20.003	TCACCCTGCAAAGGACCAAA	549
Hu.DMD.Exon55.20.004	CTCACTCACCTGCAAAGGA	550
Hu.DMD.Exon55.20.005	TCTCGCTCACTCACCTGCA	551
Hu.DMD.Exon55.20.006	CAGCCTCTCGCTCACTCACC	552
Hu.DMD.Exon55.20.007	CAAAGCAGCCTCTCGCTCAC	553
Hu.DMD.Exon55.20.008	TCTATGAGTTTCTTCCAAG	554
Hu.DMD.Exon55.20.009	GAACTGTTGCAGTAATCTAT	555
Hu.DMD.Exon55.20.010	TTCCAGGTCCAGGGGAACT	556
Hu.DMD.Exon55.20.011	CCAGGCAAGAACTTTTCCA	557
Hu.DMD.Exon55.20.012	TTCAGCTTCTGTAAGCCAGG	558
Hu.DMD.Exon55.20.013	GACATTGGCAGTTGTTTCAG	559

US 2010/0130591 A1

May 27, 2010

41

-continued

Name	Sequences	SEQ ID NO.
Hu.DMD.Exon55.20.014	GGTAGCATCCTGTAGGACAT	560
Hu.DMD.Exon55.20.015	GAGCCTTTCCTTACGGGTAG	561
Hu.DMD.Exon55.20.016	CTTGAGTCTTCTAGGAGCC	562
Hu.DMD.Exon55.20.017	CAGCTCTTTTACTCCCTTGG	563
Hu.DMD.Exon55.20.018	TTGCCATTGTTTCATCAGCT	564
Hu.DMD.Exon55.20.019	CTTACTTGCCATTGTTTCAT	565
Hu.DMD.Exon55.20.020	CCTGACTTACTTGCCATTGT	566
Hu.DMD.Exon55.20.021	AAATGCCTGACTTACTTGCC	567
Hu.DMD.Exon55.20.022	AGCGGAAATGCCTGACTTAC	568
Hu.DMD.Exon55.20.023	GCTAAAGCGGAAATGCCTGA	569
H50A(+02+30) -AVI-5656	CCACTCAGAGCTCAGATCTTCTAACTTCC	584
H50D(+07-18) -AVI-5915	GGGATCCAGTATACTTACAGGCTCC	585
H50A(+07+33)	CTTCCACTCAGAGCTCAGATCTTCTAA	586
H51A(+61+90) -AVI-4657	ACATCAAGGAAGATGGCATTCTAGTTT GG	587
H51A(+66+95) -AVI-4658	CTCCAACATCAAGGAAGATGGCATTCT AG	588
H51A(+111+134)	TTCTGTCCAAGCCCGTTGAAATC	589
H51A(+175+195)	CACCCACCATCACCCCTCYGTG	590
H51A(+199+220)	ATCATCTCGTTGATATCCTCAA	591
H51A(+66+90)	ACATCAAGGAAGATGGCATTCTAG	592
H51A(-01+25)	ACCAGAGTAACAGTCTGAGTAGGAGC	593
h51AON1	TCAAGGAAGATGGCATTCT	594
h51AON2	CCTCTGTGATTTTATAACTTGAT	595
H51D(+08-17)	ATCATTTTTTCTCATACCTTCTGCT	596
H51D(+16-07)	CTCATACCTTCTGCTTGATGATC	597
hAON#23	TGGCATTCTAGTTTGG	598
hAON#24	CCAGAGCAGGTACCTCCAACATC	599
H44A(+61+84)	TGTTTCTGCTTCTGTAGCCACTGA	600
H44A(+85+104)	TTTGTGTCTTTCTGAGAAAC	601
h44AON1	CGCCGCCATTTCTCAACAG	602
H44A(-06+14)	ATCTGTCAAATCGCCTGCAG	603
H45A(+71+90)	TGTTTTTGAGGATTGCTGAA	604
h45AON1	GCTGAATTATTTCTTCCCC	605
h45AON5	GCCCAATGCCATCCTGG	606
H45A(-06+20)	CCAATGCCATCCTGGAGTTCCTGTAA	607
H53A(+39+69)	CATTCAACTGTTGCCTCCGGTTCTGAAG GTG	608

US 2010/0130591 A1

May 27, 2010

42

-continued

Name	Sequences	SEQ ID NO.
H53A(+23+47)	CTGAAGGTGTTCTTGTACTTCATCC	609
h53AON1	CTGTTGCCTCCGGTTCTG	610
H53A(-12+10)	ATTCTTTCAACTAGATAAAAAG	611
huEx45.30.66	GCCATCCTGGAGTTCCTGTAAGATACC AAA	612
HuEx45.30.71	CCAATGCCATCCTGGAGTTCCTGTAAG ATA	613
huEx45.30.79	GCCGCTGCCC AATGCCATCCTGGAGTT CCT	614
huEx45.30.83	GTTTGCCGCTGCCC AATGCCATCCTGG AGT	615
huEx45.30.88	CAACAGTTTGCCGCTGCCCAATGCCAT CCT	616
huEx45.30.92	CTGACAACAGTTTGCCGCTGCCCAATG CCA	617
huEx45.30.96	TGTTCTGACAACAGTTTGCCGCTGCCC AAT	618
huEx45.30.99	CAATGTTCTGACAACAGTTTGCCGCTG CCC	619
huEx45.30.103	CATTCAATGTTCTGACAACAGTTTGCCG CT	620
huEx45.30.120	TATTTCTTCCCCAGTTGCATTCAATGTT CT	621
huEx45.30.127	GCTGAATTATTTCTTCCCCAGTTGCATT CA	622
huEx45.30.132	GGATTGCTGAATTATTTCTTCCCCAGTT GC	623
huEx45.30.137	TTTGAGGATTGCTGAATTATTTCTTCCC CA	624
huEx53.30.84	GTA CTTCATCCCACTGATTCTGAATTCT TT	625
huEx53.30.88	TCTTGACTTCATCCCACTGATTCTGAA TT	626
huEx53.30.91	TGTTCTTGACTTCATCCCACTGATTCT GA	627
huEx53.30.103	CGGTTCTGAAGGTGTTCTTGACTTCAT CC	628
huEx53.30.106	CTCCGGTTCTGAAGGTGTTCTTGACTT CA	629
huEx53.30.109	TGCCTCCGGTTCTGAAGGTGTTCTTGTA CT	630
huEx53.30.112	TGTTGCCCTCCGGTTCTGAAGGTGTTCTT GT	631
huEx53.30.115	AACTGTTGCCCTCCGGTTCTGAAGGTGT TCT	632

US 2010/0130591 A1

May 27, 2010

43

-continued

Name	Sequences	SEQ ID NO.
huEx53.30.118	TTCAACTGTTGCCTCCGGTTCTGAAGGT GT	633
h50AON1		
h50AON2		
<u>Peptide Transporters (NH<sub>2</sub> to COOH)*:</u>		
rTAT	RRRQRRKKRC	570
R <sub>9</sub> F <sub>2</sub>	RRRRRRRRFFC	571
(RAhx) <sub>4</sub> B	RAhxRAhxRAhxRAhxB	572
(RAhxR) <sub>4</sub> AhxB; (P007)	RAhxRAhxRAhxRAhxRAhxB	573
(AhxRR) <sub>4</sub> AhxB	AhxRAhxRAhxRAhxRAhxB	574
(RAhx) <sub>6</sub> B	RAhxRAhxRAhxRAhxRAhxRAhxB	575
(RAhx) <sub>8</sub> B	RAhxRAhxRAhxRAhxRAhxRAhxRAhxRAhxB	576
(RAhxR) <sub>5</sub> AhxB	RAhxRAhxRAhxRAhxRAhxRAhxRAhxRAhxB	577
(RAhxRRBR) <sub>2</sub> AhxB; (CP06062)	RAhxRRBRRAhxRRBRAhxB	578
MSP	ASSLNIA	579
<u>Cell Penetrating Peptide/Homing Peptide/PMO Conjugates (NH<sub>2</sub> to COOH and 5' to 3')</u>		
MSP-PMO	ASSLNIA-XB-	580
	GGCCAAACCTCGGCTTACCTGAAAT	636
CP06062-MSP-PMO	RXRRBRXRBR-XB-ASSLNIA-X-	581
	GGCCAAACCTCGGCTTACCTGAAAT	636
MSP-CP06062-PMO	ASSLNIA-X-RXRRBRXRBR-B-	582
	GGCCAAACCTCGGCTTACCTGAAAT	636
CP06062-PMO	RXRRBRXRBR-XB-	583
	GGCCAAACCTCGGCTTACCTGAAAT	636

\*Ahx is 6-aminohexanoic acid and B is beta-alanine.

## SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 651

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 25

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

&lt;400&gt; SEQUENCE: 1

ctgcaggtaa aagcatatgg atcaa

25

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 25

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

44

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 2

atcgctgca ggtaaaagca tatgg 25

<210> SEQ ID NO 3  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 3

gtcaaatacgctgc ctgcaggtaa aagca 25

<210> SEQ ID NO 4  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 4

gatctgtcaa atcgctgca ggtaa 25

<210> SEQ ID NO 5  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 5

caacagatct gtcaaatacgctgc ctgca 25

<210> SEQ ID NO 6  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 6

tttctcaaca gatctgtcaa atcgc 25

<210> SEQ ID NO 7  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 7

ccattttctca acagatctgt caaat 25

<210> SEQ ID NO 8  
<211> LENGTH: 25

US 2010/0130591 A1

May 27, 2010

45

-continued

---

```
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 8

ataatgaaaa cgccgccatt tctca                                25

<210> SEQ ID NO 9
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 9

aaatatcttt atatcataat gaaaa                                25

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 10

tgttagccac tgattaaata tcttt                                25

<210> SEQ ID NO 11
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 11

aaactgttca gcttctgtta gccac                                25

<210> SEQ ID NO 12
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 12

ttgtgtcttt ctgagaaact gttca                                25

<210> SEQ ID NO 13
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 13

ccaattctca ggaatttggtg tcttt                                25
```



US 2010/0130591 A1

May 27, 2010

46

-continued

---

<210> SEQ ID NO 14  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 14  
  
gtatttagca tgttcccaat tctca 25  
  
<210> SEQ ID NO 15  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 15  
  
cttaagatac catttggtatt tagca 25  
  
<210> SEQ ID NO 16  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 16  
  
cttaccttaa gataccattt gtatt 25  
  
<210> SEQ ID NO 17  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 17  
  
aaagacttac cttaagatac cattt 25  
  
<210> SEQ ID NO 18  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 18  
  
aaatcaaaga cttaccttaa gatac 25  
  
<210> SEQ ID NO 19  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 19  
  
aaaacaaatc aaagacttac cttaa 25

US 2010/0130591 A1

May 27, 2010

47

-continued

---

<210> SEQ ID NO 20  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 20  
  
tcgaaaaaac aaatcaaaga cttac 25  
  
<210> SEQ ID NO 21  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 21  
  
ctgtaagata ccaaaaaggc aaaac 25  
  
<210> SEQ ID NO 22  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 22  
  
cctgtaagat accaaaaagg caaaa 25  
  
<210> SEQ ID NO 23  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 23  
  
agttcctgta agatacaaaa aaggc 25  
  
<210> SEQ ID NO 24  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 24  
  
gagttcctgt aagatacaaa aaagg 25  
  
<210> SEQ ID NO 25  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 25

US 2010/0130591 A1

May 27, 2010

48

-continued

---

cctggagttc ctgtaagata ccaaa 25

<210> SEQ ID NO 26  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 26

tcctggagtt cctgtaagat accaa 25

<210> SEQ ID NO 27  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 27

gccatcctgg agttcctgta agata 25

<210> SEQ ID NO 28  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 28

tgccatcctg gagttcctgt aagat 25

<210> SEQ ID NO 29  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 29

ccaatgccat cctggagttc ctgta 25

<210> SEQ ID NO 30  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 30

cccaatgcca tcctggagtt cctgt 25

<210> SEQ ID NO 31  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

49

-continued

---

<400> SEQUENCE: 31

gctgcccaat gccatcctgg agttc

25

<210> SEQ ID NO 32

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 32

cgctgcccaa tgccatcctg gagtt

25

<210> SEQ ID NO 33

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 33

aacagtttgc cgctgcccaa tgcca

25

<210> SEQ ID NO 34

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 34

ctgacaacag ttgcccgtg cccaa

25

<210> SEQ ID NO 35

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 35

gttgcatcca atgttctgac aacag

25

<210> SEQ ID NO 36

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 36

gctgaattat ttcttcccca gttgc

25

<210> SEQ ID NO 37

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

50

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 37

attattttctt cccagttgc attca 25

<210> SEQ ID NO 38  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 38

ggcatctgtt tttgaggatt gctga 25

<210> SEQ ID NO 39  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 39

tttgaggatt gctgaattat ttctt 25

<210> SEQ ID NO 40  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 40

aatttttcct gtagaatact ggcat 25

<210> SEQ ID NO 41  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 41

atactggcat ctgtttttga ggatt 25

<210> SEQ ID NO 42  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 42

accgcagatt caggettecc aattt 25

<210> SEQ ID NO 43  
<211> LENGTH: 25  
<212> TYPE: DNA

```
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin
```

25

<400> SEQUENCE: 43

aatttttccct gtagaataact ggcat

<210> SEQ ID NO 44  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

25

<400> SEQUENCE: 44

ctgttttcgag acctcctgcc accgc

<210> SEQ ID NO 45  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

25

<400> SEQUENCE: 45

agatttcaggc ttccaatmt ttcct

<210> SEQ ID NO 46  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

25

<400> SEQUENCE: 46

ctctttttttc tgtctgacag ctggt

<210> SEQ ID NO 47  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

25

<400> SEQUENCE: 47

acctcctgcc accgcagatt caggc

<210> SEQ ID NO 48  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

25

<400> SEQUENCE: 48

cctacctctt tttctgtct gacag

<210> SEQ ID NO 49

US 2010/0130591 A1

May 27, 2010

52

-continued

---

```
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 49

gacagctggt tgcagacctc ctgcc                               25

<210> SEQ ID NO 50
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 50

gtcgccctac ctcttttttc tgtct                               25

<210> SEQ ID NO 51
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 51

gatctgtcgc cctacctctt ttttc                               25

<210> SEQ ID NO 52
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 52

tattagatct gtcgccctac ctctt                               25

<210> SEQ ID NO 53
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 53

attcctatta gatctgtcgc cctac                               25

<210> SEQ ID NO 54
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 54

agatacaaaa aaggcaaaac                               20
```



US 2010/0130591 A1

May 27, 2010

53

-continued

---

```
<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 55

aagataccaa aaaggcaaaa                20

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 56

cctgtaagat accaaaaaagg              20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 57

gagttcctgt aagataccaa                20

<210> SEQ ID NO 58
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 58

tcctggagtt cctgtaagat                20

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 59

tgccatcctg gagttcctgt                20

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 60
```

US 2010/0130591 A1

May 27, 2010

54

-continued

---

cccaatgccca tcttgagatt	20
 <210> SEQ ID NO 61 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 61	
cgctgccc aa tgccatcctg	20
 <210> SEQ ID NO 62 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 62	
ctgacaacag ttgccgctg	20
 <210> SEQ ID NO 63 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 63	
gttgcaattca atgttctgac	20
 <210> SEQ ID NO 64 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 64	
attatttctt cccagttgc	20
 <210> SEQ ID NO 65 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 65	
tttgaggatt gctgaattat	20
 <210> SEQ ID NO 66 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin	

US 2010/0130591 A1

May 27, 2010

55

-continued

---

<400> SEQUENCE: 66

atactggcat ctgtttttga 20

<210> SEQ ID NO 67

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 67

aatttttcct gtagaatact 20

<210> SEQ ID NO 68

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 68

agattcaggc ttcccaattt 20

<210> SEQ ID NO 69

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 69

acctcctgcc accgcagatt 20

<210> SEQ ID NO 70

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 70

gacagctggt tgcagacctc 20

<210> SEQ ID NO 71

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 71

ctcttttttc tgtctgacag 20

<210> SEQ ID NO 72

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

56

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 72

cctacctctt ttttctgtct 20

<210> SEQ ID NO 73  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 73

gtcgccctac ctcttttttc 20

<210> SEQ ID NO 74  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 74

gatctgtcgc cctacctctt 20

<210> SEQ ID NO 75  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 75

tattagatct gtcgccctac 20

<210> SEQ ID NO 76  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 76

attcctatta gatctgtcgc 20

<210> SEQ ID NO 77  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 77

gggggatttg agaaaataaa attac 25

<210> SEQ ID NO 78  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 78

atttgagaaa ataaaattac ctgtga                                     25

<210> SEQ ID NO 79
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 79

ctagcctgga gaaagaagaa taaaaa                                     25

<210> SEQ ID NO 80
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 80

agaaaataaaa attaccttga ctgtgc                                     25

<210> SEQ ID NO 81
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 81

ttcttctagc ctggagaaaag aagaa                                     25

<210> SEQ ID NO 82
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 82

ataaaaattac ctgtacttgc tcaag                                     25

<210> SEQ ID NO 83
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proprocessed human dystrophin

<400> SEQUENCE: 83

ttttgttctt ctagcctgga gaaag                                     25

<210> SEQ ID NO 84
<211> LENGTH: 25

```

US 2010/0130591 A1

May 27, 2010

58

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 84  
  
attaccttga cttgctcaag ctttt 25  
  
<210> SEQ ID NO 85  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 85  
  
tattcttttg ttcttctagc ctgga 25  
  
<210> SEQ ID NO 86  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 86  
  
cttgacttgc tcaagctttt ctttt 25  
  
<210> SEQ ID NO 87  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 87  
  
caagatattc ttttgttctt ctagc 25  
  
<210> SEQ ID NO 88  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 88  
  
cttttagttg ctgctctttt ccagg 25  
  
<210> SEQ ID NO 89  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 89  
  
ccaggttcaa gtgggatact agcaa 25

US 2010/0130591 A1

May 27, 2010

59

-continued

---

<210> SEQ ID NO 90  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 90  
  
atctctttga aattctgaca agata 25  
  
<210> SEQ ID NO 91  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 91  
  
agcaatgtta tctgcttcct ccaac 25  
  
<210> SEQ ID NO 92  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 92  
  
aacaaattca tttaaattctc ttga 25  
  
<210> SEQ ID NO 93  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 93  
  
ccaaccataa aacaaattca tttaa 25  
  
<210> SEQ ID NO 94  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 94  
  
ttctctcaac cataaaacaa attca 25  
  
<210> SEQ ID NO 95  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 95  
  
tttaaattctc tttaaatttc tgaca 25

US 2010/0130591 A1

May 27, 2010

60

-continued

---

<210> SEQ ID NO 96  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 96  
  
tgacaagata ttcttttggt cttct 25  
  
<210> SEQ ID NO 97  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 97  
  
ttcaagtggtg atactagcaa tgtta 25  
  
<210> SEQ ID NO 98  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 98  
  
agatatctct ttgttcttct agcct 25  
  
<210> SEQ ID NO 99  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 99  
  
ctgctctttt ccaggttcaa gtggg 25  
  
<210> SEQ ID NO 100  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 100  
  
ttcttttggt cttctagcct ggaga 25  
  
<210> SEQ ID NO 101  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 101



US 2010/0130591 A1

May 27, 2010

61

-continued

---

cttttctttt agttgctgct ctttt 25

<210> SEQ ID NO 102  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 102

ttgttcttct agcctggaga aagaa 25

<210> SEQ ID NO 103  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 103

cttctagcct ggagaaagaa gaata 25

<210> SEQ ID NO 104  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 104

agcctggaga aagaagaata aaatt 25

<210> SEQ ID NO 105  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 105

ctggagaaag aagaataaaa ttggt 25

<210> SEQ ID NO 106  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 106

gaaagaagaa taaaattggt 20

<210> SEQ ID NO 107  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

62

-continued

---

<400> SEQUENCE: 107

ggagaaagaa gaataaaatt

20

<210> SEQ ID NO 108

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 108

agcctggaga aagaagaata

20

<210> SEQ ID NO 109

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 109

cttctagcct ggagaaagaa

20

<210> SEQ ID NO 110

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 110

ttgttcttct agcctggaga

20

<210> SEQ ID NO 111

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 111

ttcttttggt cttctagcct

20

<210> SEQ ID NO 112

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 112

tgacaagata ttcttttggt

20

<210> SEQ ID NO 113

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

63

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 113

atctctttga aattctgaca 20

<210> SEQ ID NO 114  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 114

aacaaattca tttaaattctc 20

<210> SEQ ID NO 115  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 115

ttctctcaac cataaaacaa 20

<210> SEQ ID NO 116  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 116

agcaatgtta tctgcttcct 20

<210> SEQ ID NO 117  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 117

ttcaagtggg atactagcaa 20

<210> SEQ ID NO 118  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 118

ctgctctttt ccaggttcaa 20

<210> SEQ ID NO 119  
<211> LENGTH: 20  
<212> TYPE: DNA

US 2010/0130591 A1

May 27, 2010

64

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 119  
  
cttttctttt agttgctgct 20  
  
<210> SEQ ID NO 120  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 120  
  
cttgacttgc tcaagctttt 20  
  
<210> SEQ ID NO 121  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 121  
  
attaccttga cttgctcaag 20  
  
<210> SEQ ID NO 122  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 122  
  
ataaaattac cttgacttgc 20  
  
<210> SEQ ID NO 123  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 123  
  
agaaaataaa attaccttga 20  
  
<210> SEQ ID NO 124  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 124  
  
atttgagaaa ataaaattac 20  
  
<210> SEQ ID NO 125

US 2010/0130591 A1

May 27, 2010

65

-continued

---

```
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 125

gggggatttg agaaaataaa                20

<210> SEQ ID NO 126
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 126

ctgaaacaga caaatgcaac aacgt          25

<210> SEQ ID NO 127
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 127

agtaactgaa acagacaaat gcaac          25

<210> SEQ ID NO 128
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 128

ccaccagtaa ctgaaacaga caaat          25

<210> SEQ ID NO 129
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 129

cttctccacc agtaactgaa acaga          25

<210> SEQ ID NO 130
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 130

ggcaactctt ccaccagtaa ctgaa          25
```

US 2010/0130591 A1

May 27, 2010

66

-continued

---

<210> SEQ ID NO 131  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 131  
  
gcaggggcaa ctcttcacc agtaa 25  
  
<210> SEQ ID NO 132  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 132  
  
ctggcgagg ggcaactctt ccacc 25  
  
<210> SEQ ID NO 133  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 133  
  
tttaattggt tgagaattcc ctggc 25  
  
<210> SEQ ID NO 134  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 134  
  
ttgtttgaga attccctggc gcagg 25  
  
<210> SEQ ID NO 135  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 135  
  
gcacgggtcc tccagtttca tttaa 25  
  
<210> SEQ ID NO 136  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 136

US 2010/0130591 A1

May 27, 2010

67

-continued

---

tccagtttca ttttaattggt tgaga 25

<210> SEQ ID NO 137  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 137

gcttatggga gcacttacaa gcacg 25

<210> SEQ ID NO 138  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 138

tacaagcacg ggctctccag ttcca 25

<210> SEQ ID NO 139  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 139

agtttatctt gctcttctgg gctta 25

<210> SEQ ID NO 140  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 140

tctgcttgag cttattttca agttt 25

<210> SEQ ID NO 141  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 141

atcttgctct tctgggctta tggga 25

<210> SEQ ID NO 142  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

68

-continued

---

<400> SEQUENCE: 142

ctttatccac tggagatttg tctgc

25

<210> SEQ ID NO 143

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 143

cttattttca agtttatctt gctct

25

<210> SEQ ID NO 144

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 144

ctaaccttta tccactggag atttg

25

<210> SEQ ID NO 145

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 145

atttgtctgc ttgagcttat tttca

25

<210> SEQ ID NO 146

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 146

aatgtctaac ctttatccac tggag

25

<210> SEQ ID NO 147

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 147

tggttaatgt ctaaccttta tccac

25

<210> SEQ ID NO 148

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of



US 2010/0130591 A1

May 27, 2010

69

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 148

agagatgggtt aatgtctaac cttta 25

<210> SEQ ID NO 149  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 149

acggaagaga tggttaatgt ctaac 25

<210> SEQ ID NO 150  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 150

acagacaaat gcaacaacgt 20

<210> SEQ ID NO 151  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 151

ctgaaacaga caaatgcaac 20

<210> SEQ ID NO 152  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 152

agtaactgaa acagacaaat 20

<210> SEQ ID NO 153  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 153

ccaccagtaa ctgaaacaga 20

<210> SEQ ID NO 154  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

70

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 154

ctcttccacc agtaactgaa 20

<210> SEQ ID NO 155  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 155

ggcaactctt ccaccagtaa 20

<210> SEQ ID NO 156  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 156

ctggcgcagg ggcaactctt 20

<210> SEQ ID NO 157  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 157

ttgtttgaga attcctggc 20

<210> SEQ ID NO 158  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 158

tccagtttca tttaattgtt 20

<210> SEQ ID NO 159  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 159

tacaagcacg ggtcctccag 20

<210> SEQ ID NO 160  
<211> LENGTH: 20

US 2010/0130591 A1

May 27, 2010

71

-continued

---

```
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 160

gcttatggga gcacttacaa                20

<210> SEQ ID NO 161
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 161

atcttgctct tctgggctta                20

<210> SEQ ID NO 162
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 162

cttattttca agtttatctt                20

<210> SEQ ID NO 163
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 163

atttgtctgc ttgagcttat                20

<210> SEQ ID NO 164
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 164

ctttatccac tggagatttg                20

<210> SEQ ID NO 165
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 165

ctaaccttta tccactggag                20
```

US 2010/0130591 A1

May 27, 2010

72

-continued

---

<210> SEQ ID NO 166  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 166

aatgtctaac ctttatccac 20

<210> SEQ ID NO 167  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 167

tgggttaatgt ctaaccttta 20

<210> SEQ ID NO 168  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 168

agagatgggtt aatgtctaac 20

<210> SEQ ID NO 169  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 169

acggaagaga tgggttaatgt 20

<210> SEQ ID NO 170  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 170

ctgaaaggaa aatacatattt aaaaa 25

<210> SEQ ID NO 171  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 171

cctgaaaggaa aaatacatattt taaaa 25

US 2010/0130591 A1

May 27, 2010

73

-continued

---

<210> SEQ ID NO 172  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 172  
gaaacctgaa aggaaaatac atttt 25  
  
<210> SEQ ID NO 173  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 173  
ggaaacctga aaggaaaata cattt 25  
  
<210> SEQ ID NO 174  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 174  
ctctggaaac ctgaaaggaa aatac 25  
  
<210> SEQ ID NO 175  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 175  
gctctggaaa cctgaaagga aaata 25  
  
<210> SEQ ID NO 176  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 176  
gtaaagctct ggaaacctga aagga 25  
  
<210> SEQ ID NO 177  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 177

US 2010/0130591 A1

May 27, 2010

74

-continued

---

tcaggtaaag ctctggaaac ctgaa 25

<210> SEQ ID NO 178  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 178

ctcaggtaaa gctctggaaa cctga 25

<210> SEQ ID NO 179  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 179

gtttctcagg taaagctctg gaaac 25

<210> SEQ ID NO 180  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 180

tgtttctcag gtaaagctct ggaaa 25

<210> SEQ ID NO 181  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 181

aatttctcct tgtttctcag gtaaa 25

<210> SEQ ID NO 182  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 182

tttgagcttc aatttctcct tgttt 25

<210> SEQ ID NO 183  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

75

-continued

---

<400> SEQUENCE: 183

ttttatttga gcttcaattt ctcct 25

<210> SEQ ID NO 184

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 184

aagctgcccc aggtctttta ttga 25

<210> SEQ ID NO 185

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 185

aggtcttcaa gctttttttc aagct 25

<210> SEQ ID NO 186

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 186

ttcaagcttt ttttcaagct gcca 25

<210> SEQ ID NO 187

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 187

gatgatttaa ctgcttttca aggtc 25

<210> SEQ ID NO 188

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 188

ctgcttttca aggtcttcaa gcttt 25

<210> SEQ ID NO 189

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

76

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 189

aggagataac cacagcagca gatga 25

<210> SEQ ID NO 190  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 190

cagcagatga tttaactgct cttca 25

<210> SEQ ID NO 191  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 191

atttccaact gattcctaag aggag 25

<210> SEQ ID NO 192  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 192

cttggttttg ttggttataa atttc 25

<210> SEQ ID NO 193  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 193

caactgattc ctaataggag ataac 25

<210> SEQ ID NO 194  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 194

cttaacgtca aatggtcctt cttgg 25

<210> SEQ ID NO 195  
<211> LENGTH: 25  
<212> TYPE: DNA



US 2010/0130591 A1

May 27, 2010

77

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 195  
  
ttggttataa atttccaact gattc 25  
  
<210> SEQ ID NO 196  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 196  
  
cctaccttaa cgtcaaatgg tcctt 25  
  
<210> SEQ ID NO 197  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 197  
  
tccttcttgg ttggttgg tataa 25  
  
<210> SEQ ID NO 198  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 198  
  
agttccctac cttaacgtca aatgg 25  
  
<210> SEQ ID NO 199  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 199  
  
caaaaagtgc cctaccttaa cgtca 25  
  
<210> SEQ ID NO 200  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 200  
  
taaagcaaaa agttccctac cttaa 25  
  
<210> SEQ ID NO 201

US 2010/0130591 A1

May 27, 2010

78

-continued

---

<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 201  
  
atattttaaag caaaaagttc cctac 25  
  
<210> SEQ ID NO 202  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 202  
  
aggaaaatac atttttaaaaa 20  
  
<210> SEQ ID NO 203  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 203  
  
aaggaaaata catttttaaaa 20  
  
<210> SEQ ID NO 204  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 204  
  
cctgaaagga aaatacattt 20  
  
<210> SEQ ID NO 205  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 205  
  
ggaaacctga aaggaaaata 20  
  
<210> SEQ ID NO 206  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 206  
  
gctctggaaa cctgaaagga 20

US 2010/0130591 A1

May 27, 2010

79

-continued

---

<210> SEQ ID NO 207  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 207  
  
gtaaagctct ggaaacctga 20  
  
<210> SEQ ID NO 208  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 208  
  
ctcaggtaaa gctctggaaa 20  
  
<210> SEQ ID NO 209  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 209  
  
aatattctcct tgtttctcag 20  
  
<210> SEQ ID NO 210  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 210  
  
ttttatttga gcttcaattt 20  
  
<210> SEQ ID NO 211  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 211  
  
aagctgccca aggtctttta 20  
  
<210> SEQ ID NO 212  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 212

US 2010/0130591 A1

May 27, 2010

80

-continued

---

ttcaagcttt ttttcaagct	20
 <210> SEQ ID NO 213 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 213	
ctgctcttca aggtcttcaa	20
 <210> SEQ ID NO 214 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 214	
cagcagatga tttaactgct	20
 <210> SEQ ID NO 215 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 215	
aggagataac cacagcagca	20
 <210> SEQ ID NO 216 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 216	
caactgattc ctaataggag	20
 <210> SEQ ID NO 217 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 217	
ttggttataa atttccaact	20
 <210> SEQ ID NO 218 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin	

US 2010/0130591 A1

May 27, 2010

81

-continued

---

<400> SEQUENCE: 218

tccttcttgg ttggttggt

20

<210> SEQ ID NO 219

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 219

cttaacgtca aatggtcctt

20

<210> SEQ ID NO 220

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 220

cctaccttaa cgtcaaatgg

20

<210> SEQ ID NO 221

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 221

agttccctac cttaacgtca

20

<210> SEQ ID NO 222

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 222

caaaaagtgc cctaccttaa

20

<210> SEQ ID NO 223

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 223

taaagcaaaa agttccctac

20

<210> SEQ ID NO 224

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

82

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 224

atattttaaag caaaaagtgc 20

<210> SEQ ID NO 225  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 225

ctggggaaaa gaacccatat agtgc 25

<210> SEQ ID NO 226  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 226

tcctggggaa aagaacccat atagt 25

<210> SEQ ID NO 227  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 227

gtttcctggg gaaaagaacc catat 25

<210> SEQ ID NO 228  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 228

cagtttcctg gggaaaagaa cccat 25

<210> SEQ ID NO 229  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 229

tttcagtttc ctggggaaaa gaacc 25

<210> SEQ ID NO 230  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

83

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 230

tattttcagtt tcctggggaa aagaa 25

<210> SEQ ID NO 231  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 231

tgctatttca gtttctctggg gaaaa 25

<210> SEQ ID NO 232  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 232

actgctatatt cagtttctctg gggaa 25

<210> SEQ ID NO 233  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 233

tgaactgcta tttcagtttc ctggg 25

<210> SEQ ID NO 234  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 234

cttgaactgc tatttcagtt tcctg 25

<210> SEQ ID NO 235  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 235

tagcttgaac tgctatttca gtttc 25

<210> SEQ ID NO 236  
<211> LENGTH: 25

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 236

tttagcttga actgctattt cagtt                                     25

<210> SEQ ID NO 237
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 237

ttccacatcc ggttgtttag ctgta                                     25

<210> SEQ ID NO 238
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 238

tgccctttag acaaaatctc ttcca                                     25

<210> SEQ ID NO 239
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 239

tttagacaaa atctcttcca catcc                                     25

<210> SEQ ID NO 240
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 240

gtttttcctt gtacaaatgc tgccc                                     25

<210> SEQ ID NO 241
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
proporessed human dystrophin

<400> SEQUENCE: 241

gtacaaatgc tqccctttag acaaa                                     25

```



US 2010/0130591 A1

May 27, 2010

85

-continued

---

<210> SEQ ID NO 242  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 242  
  
cttcactggc tgagtggctg gtttt 25  
  
<210> SEQ ID NO 243  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 243  
  
ggctgggttt tcctgtaca aatgc 25  
  
<210> SEQ ID NO 244  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 244  
  
attaccttca ctggctgagt ggctg 25  
  
<210> SEQ ID NO 245  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 245  
  
gcttcattac cttcactggc tgagt 25  
  
<210> SEQ ID NO 246  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 246  
  
agggtgcttc attaccttca ctggc 25  
  
<210> SEQ ID NO 247  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 247  
  
gctagagggtt gcttcattac cttca 25

US 2010/0130591 A1

May 27, 2010

86

-continued

---

<210> SEQ ID NO 248  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 248  
  
atattgctag aggttgcttc attac 25  
  
<210> SEQ ID NO 249  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 249  
  
gaaaagaacc catatagtgc 20  
  
<210> SEQ ID NO 250  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 250  
  
gggaaaagaa cccatatagt 20  
  
<210> SEQ ID NO 251  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 251  
  
tcctggggaa aagaacccat 20  
  
<210> SEQ ID NO 252  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 252  
  
cagtttctctg gggaaaagaa 20  
  
<210> SEQ ID NO 253  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 253

US 2010/0130591 A1

May 27, 2010

87

-continued

---

tatttcagtt tcttggggaa 20

<210> SEQ ID NO 254  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 254

actgctatct cagtttctctg 20

<210> SEQ ID NO 255  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 255

cttgaactgc tatttcagtt 20

<210> SEQ ID NO 256  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 256

tttagcttga actgctatct 20

<210> SEQ ID NO 257  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 257

ttccacatcc ggttgcttag 20

<210> SEQ ID NO 258  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 258

tttagacaaa atctcttcca 20

<210> SEQ ID NO 259  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

88

-continued

---

<400> SEQUENCE: 259

gtacaaatgc tgccctttag

20

<210> SEQ ID NO 260

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 260

ggctggtttt tccttgtaga

20

<210> SEQ ID NO 261

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 261

cttcactggc tgagtggctg

20

<210> SEQ ID NO 262

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 262

attaccttca ctggctgagt

20

<210> SEQ ID NO 263

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 263

gcttcattac cttcactggc

20

<210> SEQ ID NO 264

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 264

agggttgcttc attaccttca

20

<210> SEQ ID NO 265

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

89

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 265

gctagaggtt gcttcattac 20

<210> SEQ ID NO 266  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 266

atattgctag aggttgcttc 20

<210> SEQ ID NO 267  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 267

ctttaacaga aaagcataca catta 25

<210> SEQ ID NO 268  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 268

tcctctttaa cagaaaagca tacac 25

<210> SEQ ID NO 269  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 269

ttcctcttta acagaaaagc ataca 25

<210> SEQ ID NO 270  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 270

taacttcctc tttaacagaa aagca 25

<210> SEQ ID NO 271  
<211> LENGTH: 25  
<212> TYPE: DNA

US 2010/0130591 A1

May 27, 2010

90

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 271  
  
ctaacttcct cttaacaga aaagc 25  
  
<210> SEQ ID NO 272  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 272  
  
tcttctaact tcctctttaa cagaa 25  
  
<210> SEQ ID NO 273  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 273  
  
atcttctaac ttctctttaa acaga 25  
  
<210> SEQ ID NO 274  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 274  
  
tcagatcttc taacttcttc tttaa 25  
  
<210> SEQ ID NO 275  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 275  
  
ctcagatctt ctaacttcct ctta 25  
  
<210> SEQ ID NO 276  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 276  
  
agagctcaga tcttctaact tcctc 25  
  
<210> SEQ ID NO 277

US 2010/0130591 A1

May 27, 2010

91

-continued

---

<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 277  
  
cagagctcag atcttctaac ttcct 25  
  
<210> SEQ ID NO 278  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 278  
  
cactcagagc tcagatcttc tact 24  
  
<210> SEQ ID NO 279  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 279  
  
ccttcactc agagctcaga tcttc 25  
  
<210> SEQ ID NO 280  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 280  
  
gtaaacggtt taccgccttc cactc 25  
  
<210> SEQ ID NO 281  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 281  
  
ctttgcctc agctcttgaa gtaaa 25  
  
<210> SEQ ID NO 282  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 282  
  
ccctcagctc ttgaagtaaa cgggt 25

US 2010/0130591 A1

May 27, 2010

92

-continued

---

<210> SEQ ID NO 283  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 283  
  
ccaggagcta ggtcaggctg ctttg 25  
  
<210> SEQ ID NO 284  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 284  
  
ggtcaggctg ctttgccctc agctc 25  
  
<210> SEQ ID NO 285  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 285  
  
aggctccaat agtggtcagt ccagg 25  
  
<210> SEQ ID NO 286  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 286  
  
tcagtcagg agctaggtca ggctg 25  
  
<210> SEQ ID NO 287  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 287  
  
cttacaggct ccaatagtgg tcagt 25  
  
<210> SEQ ID NO 288  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 288



US 2010/0130591 A1

May 27, 2010

93

-continued

---

gtatacttac aggtccaat agtgg 25

<210> SEQ ID NO 289  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 289

atccagtata cttacaggct ccaat 25

<210> SEQ ID NO 290  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 290

atgggatcca gtatacttac aggct 25

<210> SEQ ID NO 291  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 291

agagaatggg atccagtata cttac 25

<210> SEQ ID NO 292  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 292

acagaaaagc atacacatta 20

<210> SEQ ID NO 293  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 293

tttaacagaa aagcatacac 20

<210> SEQ ID NO 294  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

94

-continued

---

<400> SEQUENCE: 294

tcctctttaa cagaaaagca

20

<210> SEQ ID NO 295

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 295

taacttcctc tttaacagaa

20

<210> SEQ ID NO 296

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 296

tcttctaact tcctctttaa

20

<210> SEQ ID NO 297

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 297

tcagatcttc taacttcctc

20

<210> SEQ ID NO 298

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 298

ccttcactc agagctcaga

20

<210> SEQ ID NO 299

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 299

gtaaacgggt taccgccttc

20

<210> SEQ ID NO 300

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

95

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 300

ccctcagctc ttgaagtaaa 20

<210> SEQ ID NO 301  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 301

ggtcaggctg ctttgccctc 20

<210> SEQ ID NO 302  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 302

tcagtcagg agctaggta 20

<210> SEQ ID NO 303  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 303

aggctccaat agtggtcagt 20

<210> SEQ ID NO 304  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 304

cttacaggct ccaatagtgg 20

<210> SEQ ID NO 305  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 305

gtatacttac aggctccaat 20

<210> SEQ ID NO 306  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

96

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 306

atccagtata cttacaggct 20

<210> SEQ ID NO 307  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 307

atgggatcca gtatacttac 20

<210> SEQ ID NO 308  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 308

agagaatggg atccagtata 20

<210> SEQ ID NO 309  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 309

ctaaaatatt ttgggttttt gcaaaa 26

<210> SEQ ID NO 310  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 310

gctaaaatat ttgggttttt tgcaaa 26

<210> SEQ ID NO 311  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 311

taggagctaa aatatttttg gttttt 26

<210> SEQ ID NO 312  
<211> LENGTH: 25

US 2010/0130591 A1

May 27, 2010

97

-continued

---

```
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 312

agtaggagct aaaatatttt ggggtt                25

<210> SEQ ID NO 313
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 313

tgagtaggag ctaaaatatt ttggg                25

<210> SEQ ID NO 314
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 314

ctgagtagga gctaaaatat ttggg                26

<210> SEQ ID NO 315
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 315

cagtctgagt aggagctaaa atatt                25

<210> SEQ ID NO 316
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 316

acagtctgag taggagctaa aatatt                26

<210> SEQ ID NO 317
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 317

gagtaacagt ctgagtagga gctaaa                26
```

US 2010/0130591 A1

May 27, 2010

98

-continued

---

<210> SEQ ID NO 318  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 318  
  
cagagtaaca gtctgagtag gagct 25  
  
<210> SEQ ID NO 319  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 319  
  
caccagagta acagtctgag taggag 26  
  
<210> SEQ ID NO 320  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 320  
  
gtcaccagag taacagtctg agtag 25  
  
<210> SEQ ID NO 321  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 321  
  
aaccacaggt tgtgtcacca gagtaa 26  
  
<210> SEQ ID NO 322  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 322  
  
gttgtgtcac cagagtaaca gtctg 25  
  
<210> SEQ ID NO 323  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 323  
  
tggcagtttc cttagtaacc acaggt 26

US 2010/0130591 A1

May 27, 2010

99

-continued

---

<210> SEQ ID NO 324  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 324  
  
atttctagtt tggagatggc agtttc 26  
  
<210> SEQ ID NO 325  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 325  
  
ggaagatggc atttctagtt tggag 25  
  
<210> SEQ ID NO 326  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 326  
  
catcaaggaa gatggcattt ctagtt 26  
  
<210> SEQ ID NO 327  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 327  
  
gagcaggtac ctccaacatc aaggaa 26  
  
<210> SEQ ID NO 328  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 328  
  
atctgccaga gcaggtacct ccaac 25  
  
<210> SEQ ID NO 329  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 329

US 2010/0130591 A1

May 27, 2010

100

-continued

---

aagttctgtc caagcccggt tgaaat 26

<210> SEQ ID NO 330  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 330

cgggtgaaat ctgccagagc aggtac 26

<210> SEQ ID NO 331  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 331

gagaaagcca gtcggtaagt tctgtc 26

<210> SEQ ID NO 332  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 332

gtcggtaagt tctgtccaag cccgg 25

<210> SEQ ID NO 333  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 333

ataacttgat caagcagaga aagcca 26

<210> SEQ ID NO 334  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 334

aagcagagaa agccagtcgg taagt 25

<210> SEQ ID NO 335  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin



US 2010/0130591 A1

May 27, 2010

101

-continued

---

<400> SEQUENCE: 335

caccctctgt gattttataa cttgat

26

<210> SEQ ID NO 336

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 336

caaggtcacc caccatcacc ctctgt

26

<210> SEQ ID NO 337

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 337

catcacctc tgtgatttta taact

25

<210> SEQ ID NO 338

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 338

cttctgcttg atgatcatct cgttga

26

<210> SEQ ID NO 339

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 339

ccttctgctt gatgatcatc tcgttg

26

<210> SEQ ID NO 340

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 340

atctcgttga taccctcaag gtcacc

26

<210> SEQ ID NO 341

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

102

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 341

tcataccttc tgcttgatga tcattct 26

<210> SEQ ID NO 342  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 342

tcatttttttc tcataccttc tgcttg 26

<210> SEQ ID NO 343  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 343

tttttcata ccttctgctt gatgat 26

<210> SEQ ID NO 344  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 344

ttttatcatt ttttctcata ccttct 26

<210> SEQ ID NO 345  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 345

ccaaacttta tcatttttttc tcatac 26

<210> SEQ ID NO 346  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 346

atattttggg tttttgcaaa 20

<210> SEQ ID NO 347  
<211> LENGTH: 20  
<212> TYPE: DNA

US 2010/0130591 A1

May 27, 2010

103

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 347  
  
aaaaatatttt gggtttttgc 20  
  
<210> SEQ ID NO 348  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 348  
  
gagctaaaat attttgggtt 20  
  
<210> SEQ ID NO 349  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 349  
  
agtaggagct aaaatatttt 20  
  
<210> SEQ ID NO 350  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 350  
  
gtctgagtag gagctaaaat 20  
  
<210> SEQ ID NO 351  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 351  
  
taacagtctg agtaggagct 20  
  
<210> SEQ ID NO 352  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 352  
  
cagagtaaca gtctgagtag 20  
  
<210> SEQ ID NO 353

US 2010/0130591 A1

May 27, 2010

104

-continued

---

<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 353  
  
cacaggttgt gtcaccagag 20  
  
<210> SEQ ID NO 354  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 354  
  
agtttcctta gtaaccacag 20  
  
<210> SEQ ID NO 355  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 355  
  
tagtttggag atggcagttt 20  
  
<210> SEQ ID NO 356  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 356  
  
ggaagatggc atttctagtt 20  
  
<210> SEQ ID NO 357  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 357  
  
tacctccaac atcaaggaag 20  
  
<210> SEQ ID NO 358  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 358  
  
atctgccaga gcaggtacct 20

US 2010/0130591 A1

May 27, 2010

105

-continued

---

<210> SEQ ID NO 359  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 359  
  
ccaagcccggttgaaatctg 20  
  
<210> SEQ ID NO 360  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 360  
  
gtcggtaagt tctgtccaag 20  
  
<210> SEQ ID NO 361  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 361  
  
aagcagagaa agccagtcgg 20  
  
<210> SEQ ID NO 362  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 362  
  
ttttataact tgatcaagca 20  
  
<210> SEQ ID NO 363  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 363  
  
catcacctc tgtgatttta 20  
  
<210> SEQ ID NO 364  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 364

US 2010/0130591 A1

May 27, 2010

106

-continued

---

ctcaagggtca cccaccatca 20

<210> SEQ ID NO 365  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 365

catctcgttg atatcctcaa 20

<210> SEQ ID NO 366  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 366

cttctgcttg atgatcatct 20

<210> SEQ ID NO 367  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 367

cataccttct gcttgatgat 20

<210> SEQ ID NO 368  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 368

tttctcatatc cttctgcttg 20

<210> SEQ ID NO 369  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 369

catTTTTTct cataccttct 20

<210> SEQ ID NO 370  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

107

-continued

---

<400> SEQUENCE: 370

tttatcattt tttctcatac

20

<210> SEQ ID NO 371

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 371

caacttttat cattttttct

20

<210> SEQ ID NO 372

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 372

ctgtaagaac aaatatccct tagta

25

<210> SEQ ID NO 373

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 373

tgccctgtaag aacaaatata cctta

25

<210> SEQ ID NO 374

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 374

gttgccctgta agaacaaata tccct

25

<210> SEQ ID NO 375

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 375

attgttgcct gtaagaacaa atatac

25

<210> SEQ ID NO 376

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

108

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 376

gcattgttgc ctgtaagaac aaata 25

<210> SEQ ID NO 377  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 377

cctgcattgt tgcctgtaag aacaa 25

<210> SEQ ID NO 378  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 378

atcctgcatt gttgcctgta agaac 25

<210> SEQ ID NO 379  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 379

caaatcctgc attgttgcct gtaag 25

<210> SEQ ID NO 380  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 380

tccaaatcct gcattgttgc ctgta 25

<210> SEQ ID NO 381  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 381

tgttccaaat cctgcattgt tgcct 25

<210> SEQ ID NO 382  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence



US 2010/0130591 A1

May 27, 2010

109

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 382

tctgttccaa atcctgcatt gttgc 25

<210> SEQ ID NO 383  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 383

aactggggac gcctctgttc caaat 25

<210> SEQ ID NO 384  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 384

gcctctgttc caaatcctgc attgt 25

<210> SEQ ID NO 385  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 385

cagcggtaat gagttcttcc aactg 25

<210> SEQ ID NO 386  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 386

cttccaactg gggacgcctc tgttc 25

<210> SEQ ID NO 387  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 387

cttggttttc aaattttggg cagcg 25

<210> SEQ ID NO 388  
<211> LENGTH: 25

US 2010/0130591 A1

May 27, 2010

110

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 388  
  
ctagcctctt gattgctggt cttgt 25  
  
<210> SEQ ID NO 389  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 389  
  
ttttcaaatt ttgggcagcg gtaat 25  
  
<210> SEQ ID NO 390  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 390  
  
ttcgatccgt aatgattgtt ctage 25  
  
<210> SEQ ID NO 391  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 391  
  
gattgctggt cttgtttttc aaatt 25  
  
<210> SEQ ID NO 392  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 392  
  
cttacttcga tccgtaatga ttgtt 25  
  
<210> SEQ ID NO 393  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 393  
  
ttgttctagc ctcttgattg ctggt 25

US 2010/0130591 A1

May 27, 2010

111

-continued

---

<210> SEQ ID NO 394  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 394  
  
aaaaacttac ttcgatccgt aatga 25  
  
<210> SEQ ID NO 395  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 395  
  
tgttaaaaaa cttacttcga tccgt 25  
  
<210> SEQ ID NO 396  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 396  
  
atgcttggtta aaaaacttac ttcga 25  
  
<210> SEQ ID NO 397  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 397  
  
gtcccatgct tgttaaaaaa cttac 25  
  
<210> SEQ ID NO 398  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 398  
  
agaacaaata tcccttagta 20  
  
<210> SEQ ID NO 399  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 399  
  
gtaagaacaa atatccctta 20

US 2010/0130591 A1

May 27, 2010

112

-continued

---

<210> SEQ ID NO 400  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 400  
tgccctgtaag aacaaatattc 20  
  
<210> SEQ ID NO 401  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 401  
attgttgccct gtaagaacaa 20  
  
<210> SEQ ID NO 402  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 402  
cctgcattgt tgccctgtaag 20  
  
<210> SEQ ID NO 403  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 403  
caaatcctgc attgttgccct 20  
  
<210> SEQ ID NO 404  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 404  
gcctctgttc caaatcctgc 20  
  
<210> SEQ ID NO 405  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 405

US 2010/0130591 A1

May 27, 2010

113

-continued

---

cttccaactg gggacgcctc 20

<210> SEQ ID NO 406  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 406

cagcggtaat gagttcttcc 20

<210> SEQ ID NO 407  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 407

ttttcaaatt ttgggcagcg 20

<210> SEQ ID NO 408  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 408

gattgctggt cttgttttcc 20

<210> SEQ ID NO 409  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 409

ttgttctagc ctcttgattg 20

<210> SEQ ID NO 410  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 410

ttcgatccgt aatgattgtt 20

<210> SEQ ID NO 411  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

114

-continued

---

<400> SEQUENCE: 411  
cttacttcga tccgtaatga 20

<210> SEQ ID NO 412  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 412  
aaaaaacttac ttcgatccgt 20

<210> SEQ ID NO 413  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 413  
tgttaaaaaa cttacttcga 20

<210> SEQ ID NO 414  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 414  
atgcttggtta aaaaacttac 20

<210> SEQ ID NO 415  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 415  
gtcccatgct tgttaaaaaa 20

<210> SEQ ID NO 416  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 416  
ctagaataaa aggaaaaata aatat 25

<210> SEQ ID NO 417  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

115

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 417

aactagaata aaaggaaaaa taaat 25

<210> SEQ ID NO 418

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 418

ttcaactaga ataaaaggaa aaata 25

<210> SEQ ID NO 419

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 419

ctttcaacta gaataaaaagg aaaaa 25

<210> SEQ ID NO 420

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 420

attctttcaa ctagaataaa aggaa 25

<210> SEQ ID NO 421

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 421

gaattctttc aactagaata aaagg 25

<210> SEQ ID NO 422

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 422

tctgaattct ttcaactaga ataaa 25

<210> SEQ ID NO 423

<211> LENGTH: 25

<212> TYPE: DNA

US 2010/0130591 A1

May 27, 2010

116

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 423  
  
attctgaatt ctttcaacta gaata 25  
  
<210> SEQ ID NO 424  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 424  
  
ctgattctga attctttcaa ctaga 25  
  
<210> SEQ ID NO 425  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 425  
  
cactgattct gaattctttc aacta 25  
  
<210> SEQ ID NO 426  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 426  
  
tcccactgat tctgaattct ttcaa 25  
  
<210> SEQ ID NO 427  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 427  
  
catccactg attctgaatt ctttc 25  
  
<210> SEQ ID NO 428  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 428  
  
tacttcattcc cactgattct gaatt 25  
  
<210> SEQ ID NO 429



US 2010/0130591 A1

May 27, 2010

117

-continued

---

<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 429  
  
ctgaaggtgt tcttgtactt catcc 25  
  
<210> SEQ ID NO 430  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 430  
  
cggttctgaa ggtgttcttg tact 24  
  
<210> SEQ ID NO 431  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 431  
  
ctgttgccctc cggttctgaa ggtgt 25  
  
<210> SEQ ID NO 432  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 432  
  
tttcattcaa ctgttgccctc cggtt 25  
  
<210> SEQ ID NO 433  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 433  
  
taacatttca ttcaactggt gacct 25  
  
<210> SEQ ID NO 434  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 434  
  
ttgtgttgaa tcctttaaca tttca 25

US 2010/0130591 A1

May 27, 2010

118

-continued

---

<210> SEQ ID NO 435  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 435  
  
tcttccttag cttccagcca ttgtg 25  
  
<210> SEQ ID NO 436  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 436  
  
cttagcttcc agccattgtg ttgaa 25  
  
<210> SEQ ID NO 437  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 437  
  
gtcctaagac ctgctcagct tcttc 25  
  
<210> SEQ ID NO 438  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 438  
  
ctgctcagct tcttccttag cttcc 25  
  
<210> SEQ ID NO 439  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 439  
  
ctcaagcttg gctctggcct gtcct 25  
  
<210> SEQ ID NO 440  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 440

US 2010/0130591 A1

May 27, 2010

119

-continued

---

ggcctgtcct aagacctgct cagct 25

<210> SEQ ID NO 441  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 441

tagggaccct ccttccatga ctcaa 25

<210> SEQ ID NO 442  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 442

tttggttg atctactgta taggg 25

<210> SEQ ID NO 443  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 443

accctccttc catgactcaa gcttg 25

<210> SEQ ID NO 444  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 444

cttggtttct gtgattttct ttgg 25

<210> SEQ ID NO 445  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 445

atctactgta tagggaccct ccttc 25

<210> SEQ ID NO 446  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

120

-continued

---

<400> SEQUENCE: 446

ctaaccttgg tttctgtgat tttct 25

<210> SEQ ID NO 447

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 447

tttcttttgg attgcatcta ctgta 25

<210> SEQ ID NO 448

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 448

tgataactaac cttggtttct gtgat 25

<210> SEQ ID NO 449

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 449

atctttgata ctaaccttgg tttct 25

<210> SEQ ID NO 450

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 450

aaggatatctt tgataactaac cttgg 25

<210> SEQ ID NO 451

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 451

ttaaaaagggt atctttgata ctaac 25

<210> SEQ ID NO 452

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

121

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 452

ataaaaggaa aaataaatat 20

<210> SEQ ID NO 453  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 453

gaataaaaagg aaaaataaat 20

<210> SEQ ID NO 454  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 454

aactagaata aaaggaaaaa 20

<210> SEQ ID NO 455  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 455

ctttcaacta gaataaaaagg 20

<210> SEQ ID NO 456  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 456

gaattctttc aactagaata 20

<210> SEQ ID NO 457  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 457

attctgaatt ctttcaacta 20

<210> SEQ ID NO 458  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

122

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 458

tacttcatcc cactgattct 20

<210> SEQ ID NO 459  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 459

ctgaagggtgt tcttgatct 19

<210> SEQ ID NO 460  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 460

ctgttgccctc cggttctgaa 20

<210> SEQ ID NO 461  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 461

taacatttca ttcaactgtt 20

<210> SEQ ID NO 462  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 462

ttgtgttgaa tcctttaaca 20

<210> SEQ ID NO 463  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 463

cttagcttcc agccattgtg 20

<210> SEQ ID NO 464  
<211> LENGTH: 20

US 2010/0130591 A1

May 27, 2010

123

-continued

---

```
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 464

ctgctcagct tcttccttag                20

<210> SEQ ID NO 465
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 465

ggcctgtcct aagacctgct                20

<210> SEQ ID NO 466
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 466

ctcaagcttg gctctggcct                20

<210> SEQ ID NO 467
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 467

accctccttc catgactcaa                20

<210> SEQ ID NO 468
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 468

atctactgta tagggaccct                20

<210> SEQ ID NO 469
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
      proprocessed human dystrophin

<400> SEQUENCE: 469

tttcttttgg attgcatcta                20
```

US 2010/0130591 A1

May 27, 2010

124

-continued

---

<210> SEQ ID NO 470  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 470  
  
cttggtttct gtgattttct 20  
  
<210> SEQ ID NO 471  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 471  
  
ctaaccttgg tttctgtgat 20  
  
<210> SEQ ID NO 472  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 472  
  
tgataactaac cttggtttct 20  
  
<210> SEQ ID NO 473  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 473  
  
atctttgata ctaaccttgg 20  
  
<210> SEQ ID NO 474  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 474  
  
aaggatatctt tgataactaac 20  
  
<210> SEQ ID NO 475  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 475  
  
ttaaaaaggt atctttgata 20



US 2010/0130591 A1

May 27, 2010

125

-continued

---

<210> SEQ ID NO 476  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 476  
  
ctatagattt ttatgagaaa gaga 24  
  
<210> SEQ ID NO 477  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 477  
  
aactgctata gattttttatg agaaa 25  
  
<210> SEQ ID NO 478  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 478  
  
tggccaactg ctatagattt ttatg 25  
  
<210> SEQ ID NO 479  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 479  
  
gtctttggcc aactgctata gattt 25  
  
<210> SEQ ID NO 480  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 480  
  
cggaggtctt tggccaactg ctata 25  
  
<210> SEQ ID NO 481  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 481

US 2010/0130591 A1

May 27, 2010

126

-continued

---

actggcggag gtctttggcc aactg 25

<210> SEQ ID NO 482  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 482

tttgtctgcc actggcggag gtctt 25

<210> SEQ ID NO 483  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 483

agtcatttgc cacatctaca ttgt 25

<210> SEQ ID NO 484  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 484

tttgccacat ctacatttgt ctgcc 25

<210> SEQ ID NO 485  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 485

ccggagaagt ttcagggccca agtca 25

<210> SEQ ID NO 486  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 486

gtatcatctg cagaataatc ccgga 25

<210> SEQ ID NO 487  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

127

-continued

---

<400> SEQUENCE: 487

taatcccga gaagtttcag ggcca

25

<210> SEQ ID NO 488

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 488

ttatcatgtg gacttttctg gtatc

25

<210> SEQ ID NO 489

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 489

agaggcattg atattctctg ttatc

25

<210> SEQ ID NO 490

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 490

atgtggactt ttctggtatc atctg

25

<210> SEQ ID NO 491

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 491

cttttatgaa tgcttctcca agagg

25

<210> SEQ ID NO 492

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 492

atattctctg ttatcatgtg gactt

25

<210> SEQ ID NO 493

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

128

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 493

catacctttt atgaatgctt ctcca 25

<210> SEQ ID NO 494

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 494

ctccaagagg cattgatatt ctctg 25

<210> SEQ ID NO 495

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 495

taattcatac cttttatgaa tgctt 25

<210> SEQ ID NO 496

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 496

cttttatgaa tgcttctcca agagg 25

<210> SEQ ID NO 497

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 497

taatgtaatt catacctttt atgaa 25

<210> SEQ ID NO 498

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 498

agaaataatg taattcatac ctttt 25

<210> SEQ ID NO 499

<211> LENGTH: 25

<212> TYPE: DNA

US 2010/0130591 A1

May 27, 2010

129

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 499  
  
gttttagaaa taatgtaatt catac 25  
  
<210> SEQ ID NO 500  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 500  
  
gatttttatg agaaagaga 19  
  
<210> SEQ ID NO 501  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 501  
  
ctatagattt ttatgagaaa 20  
  
<210> SEQ ID NO 502  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 502  
  
aactgctata gatttttatg 20  
  
<210> SEQ ID NO 503  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 503  
  
tggccaactg ctatagattt 20  
  
<210> SEQ ID NO 504  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 504  
  
gtctttggcc aactgctata 20  
  
<210> SEQ ID NO 505

US 2010/0130591 A1

May 27, 2010

130

-continued

---

<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 505  
  
cggaggtctt tggccaactg 20  
  
<210> SEQ ID NO 506  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 506  
  
tttgtctgcc actggcggag 20  
  
<210> SEQ ID NO 507  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 507  
  
tttgccacat ctacatttgt 20  
  
<210> SEQ ID NO 508  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 508  
  
ttcagggccca agtcatttgc 20  
  
<210> SEQ ID NO 509  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 509  
  
taatcccgga gaagtttcag 20  
  
<210> SEQ ID NO 510  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 510  
  
gtatcatctg cagaataatc 20

US 2010/0130591 A1

May 27, 2010

131

-continued

---

<210> SEQ ID NO 511  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 511  
  
atgtggactt ttctggtatc 20  
  
<210> SEQ ID NO 512  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 512  
  
atattctctg ttatcatgtg 20  
  
<210> SEQ ID NO 513  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 513  
  
ctccaagagg cattgatatt 20  
  
<210> SEQ ID NO 514  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 514  
  
cttttatgaa tgcttctcca 20  
  
<210> SEQ ID NO 515  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 515  
  
catacctttt atgaatgctt 20  
  
<210> SEQ ID NO 516  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 516

US 2010/0130591 A1

May 27, 2010

132

-continued

---

taattcatac cttttatgaa	20
 <210> SEQ ID NO 517 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 517	
taatgtaatt catacctttt	20
 <210> SEQ ID NO 518 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 518	
agaaataatg taattcatac	20
 <210> SEQ ID NO 519 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 519	
gttttagaaa taatgtaatt	20
 <210> SEQ ID NO 520 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 520	
ctgcaaagga ccaaagtgtc agatg	25
 <210> SEQ ID NO 521 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin  <400> SEQUENCE: 521	
tcaccctgca aaggacaaaa tgttc	25
 <210> SEQ ID NO 522 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense sequence to target splice site of proprocessed human dystrophin	



US 2010/0130591 A1

May 27, 2010

133

-continued

---

<400> SEQUENCE: 522

ctcactcacc ctgcaaagga ccaaa 25

<210> SEQ ID NO 523

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 523

tctcgctcac tcaccctgca aagga 25

<210> SEQ ID NO 524

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 524

cagcctctcg ctcactcacc ctgca 25

<210> SEQ ID NO 525

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 525

caaagcagcc tctcgctcac tcacc 25

<210> SEQ ID NO 526

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 526

tcttccaaag cagcctctcg ctcac 25

<210> SEQ ID NO 527

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 527

tctatgagtt tcttccaaag cagcc 25

<210> SEQ ID NO 528

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of

US 2010/0130591 A1

May 27, 2010

134

-continued

---

proprocessed human dystrophin

<400> SEQUENCE: 528

gttgcagtaa tctatgagtt tcttc 25

<210> SEQ ID NO 529  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 529

gaactgttgc agtaatctat gagtt 25

<210> SEQ ID NO 530  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 530

ttccaggtcc agggggaact gttgc 25

<210> SEQ ID NO 531  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 531

gtaagccagg caagaaactt ttcca 25

<210> SEQ ID NO 532  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 532

ccaggcaaga aactttttcca ggtcc 25

<210> SEQ ID NO 533  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 533

tggcagttgt ttcagcttct gtaag 25

<210> SEQ ID NO 534  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

135

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 534

ggtagcatcc tgtaggacat tggca 25

<210> SEQ ID NO 535  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 535

gacattggca gttgtttcag cttct 25

<210> SEQ ID NO 536  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 536

tctaggagcc tttccttacg ggtag 25

<210> SEQ ID NO 537  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 537

cttttactcc cttggagtct tctag 25

<210> SEQ ID NO 538  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 538

gagcctttcc ttacgggtag catcc 25

<210> SEQ ID NO 539  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 539

ttgccattgt ttcacagct ctttt 25

<210> SEQ ID NO 540  
<211> LENGTH: 25

US 2010/0130591 A1

May 27, 2010

136

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 540  
  
cttggagtct tctaggagcc tttcc 25  
  
<210> SEQ ID NO 541  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 541  
  
cttacttgcc attgtttcat cagct 25  
  
<210> SEQ ID NO 542  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 542  
  
cagctctttt actcccttgg agtct 25  
  
<210> SEQ ID NO 543  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 543  
  
cctgacttac ttgccattgt ttcac 25  
  
<210> SEQ ID NO 544  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 544  
  
aaatgcctga cttacttgcc attgt 25  
  
<210> SEQ ID NO 545  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 545  
  
agcggaaatg cctgacttac ttgcc 25

US 2010/0130591 A1

May 27, 2010

137

-continued

---

<210> SEQ ID NO 546  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 546  
  
gctaaagcgg aaatgcctga cttac 25  
  
<210> SEQ ID NO 547  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 547  
  
aaggacaaa tggtcagatg 20  
  
<210> SEQ ID NO 548  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 548  
  
ctgcaaagga ccaaatgttc 20  
  
<210> SEQ ID NO 549  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 549  
  
tcaccctgca aaggacaaa 20  
  
<210> SEQ ID NO 550  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 550  
  
ctcactcacc ctgcaaagga 20  
  
<210> SEQ ID NO 551  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 551  
  
tctcgtcac tcaccctgca 20

US 2010/0130591 A1

May 27, 2010

138

-continued

---

<210> SEQ ID NO 552  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 552  
cagcctctcg ctcactcacc 20  
  
<210> SEQ ID NO 553  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 553  
caaagcagcc tctcgctcac 20  
  
<210> SEQ ID NO 554  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 554  
tctatgagtt tcttccaaag 20  
  
<210> SEQ ID NO 555  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 555  
gaactgttgc agtaatctat 20  
  
<210> SEQ ID NO 556  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 556  
ttccaggtcc agggggaact 20  
  
<210> SEQ ID NO 557  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 557

US 2010/0130591 A1

May 27, 2010

139

-continued

---

ccaggcaaga aacttttcca 20

<210> SEQ ID NO 558  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 558

ttcagettct gtaagccagg 20

<210> SEQ ID NO 559  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 559

gacattggca gttgtttcag 20

<210> SEQ ID NO 560  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 560

ggtagcatcc tgtaggacat 20

<210> SEQ ID NO 561  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 561

gagcctttcc ttacgggtag 20

<210> SEQ ID NO 562  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 562

cttgaggtct tctaggagcc 20

<210> SEQ ID NO 563  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

140

-continued

---

<400> SEQUENCE: 563

cagctctttt actcccttgg

20

<210> SEQ ID NO 564

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 564

ttgccattgt ttcacagct

20

<210> SEQ ID NO 565

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 565

cttacttgcc attgtttcat

20

<210> SEQ ID NO 566

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 566

cctgacttac ttgccattgt

20

<210> SEQ ID NO 567

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 567

aaatgcctga cttacttgcc

20

<210> SEQ ID NO 568

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 568

agcggaaatg cctgacttac

20

<210> SEQ ID NO 569

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:



US 2010/0130591 A1

May 27, 2010

141

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 569

gctaaagcgg aaatgcctga

20

<210> SEQ ID NO 570

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO

<400> SEQUENCE: 570

Arg Arg Arg Gln Arg Arg Lys Lys Arg Cys

1 5 10

<210> SEQ ID NO 571

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO

<400> SEQUENCE: 571

Arg Arg Arg Arg Arg Arg Arg Arg Phe Phe Cys

1 5 10

<210> SEQ ID NO 572

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 3, 6, 9, 12

<223> OTHER INFORMATION: X = 6-aminohexanoic acid

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 13

<223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 572

Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Xaa

1 5 10

<210> SEQ ID NO 573

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 2, 5, 8, 11, 13

<223> OTHER INFORMATION: X = 6-aminohexanoic acid

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 14

<223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 573

US 2010/0130591 A1

May 27, 2010

142

-continued

---

Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Xaa Xaa  
1 5 10

<210> SEQ ID NO 574  
<211> LENGTH: 14  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 1, 4, 7, 10, 13  
<223> OTHER INFORMATION: X = 6-aminohexanoic acid  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 14  
<223> OTHER INFORMATION: X = beta alanine  
  
<400> SEQUENCE: 574

Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Xaa Xaa  
1 5 10

<210> SEQ ID NO 575  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 2, 4, 6, 8, 10, 12  
<223> OTHER INFORMATION: X = 6-aminohexanoic acid  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 13  
<223> OTHER INFORMATION: X = beta alanine  
  
<400> SEQUENCE: 575

Arg Xaa Arg Xaa Arg Xaa Arg Xaa Arg Xaa Arg Xaa Xaa  
1 5 10

<210> SEQ ID NO 576  
<211> LENGTH: 17  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 2, 4, 6, 8, 10, 12, 14, 16  
<223> OTHER INFORMATION: X = 6-aminohexanoic acid  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 17  
<223> OTHER INFORMATION: X = beta alanine  
  
<400> SEQUENCE: 576

Arg Xaa Arg Xaa Arg Xaa Arg Xaa Arg Xaa Arg Xaa Arg Xaa  
1 5 10 15

Xaa

<210> SEQ ID NO 577  
<211> LENGTH: 17  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

143

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 2, 5, 8, 11, 14, 16  
<223> OTHER INFORMATION: X = 6-aminohexanoic acid  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 17  
<223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 577

Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Xaa  
1 5 10 15

Xaa

<210> SEQ ID NO 578  
<211> LENGTH: 14  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 2, 8, 13  
<223> OTHER INFORMATION: X = 6-aminohexanoic acid  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 5, 11, 14  
<223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 578

Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Xaa Xaa  
1 5 10

<210> SEQ ID NO 579  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Peptide Transporter for Intracellular Delivery  
of PMO

<400> SEQUENCE: 579

Ala Ser Ser Leu Asn Ile Ala  
1 5

<210> SEQ ID NO 580  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Cell Penetrating Peptide / Homing Peptide /  
PMO Conjugates  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 8  
<223> OTHER INFORMATION: X = a neutral amino acid C(=O) (CHR)n NH-,  
where n is 2 to 7 and each R is independently H or methyl  
<220> FEATURE:  
<221> NAME/KEY: MOD\_RES  
<222> LOCATION: 9  
<223> OTHER INFORMATION: X = Beta alanine

<400> SEQUENCE: 580

Ala Ser Ser Leu Asn Ile Ala Xaa Xaa

US 2010/0130591 A1

May 27, 2010

144

-continued

---

1 5

<210> SEQ ID NO 581  
 <211> LENGTH: 22  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Cell Penetrating Peptide / Homing Peptide /  
 PMO Conjugates  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 2, 8, 13, 22  
 <223> OTHER INFORMATION: X = a neutral amino acid C(=O) (CHR)n NH-,  
 where n is 2 to 7 and each R is independently H or methyl  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 5, 11, 14  
 <223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 581

Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Xaa Ala Ser  
 1 5 10 15

Ser Leu Asn Ile Ala Xaa  
 20

<210> SEQ ID NO 582  
 <211> LENGTH: 21  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Cell Penetrating Peptide / Homing Peptide /  
 PMO Conjugates  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 8, 10, 16  
 <223> OTHER INFORMATION: X = a neutral amino acid C(=O) (CHR)n NH-,  
 where n is 2 to 7 and each R is independently H or methyl  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 13, 19, 21  
 <223> OTHER INFORMATION: Xaa = beta alanine

<400> SEQUENCE: 582

Ala Ser Ser Leu Asn Ile Ala Xaa Arg Xaa Arg Arg Xaa Arg Arg Xaa  
 1 5 10 15

Arg Arg Xaa Arg Xaa  
 20

<210> SEQ ID NO 583  
 <211> LENGTH: 14  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Cell Penetrating Peptide / Homing Peptide /  
 PMO Conjugates  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 2, 8, 13  
 <223> OTHER INFORMATION: X = a neutral amino acid C(=O) (CHR)n NH-,  
 where n is 2 to 7 and each R is independently H or methyl  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: 5, 11, 14  
 <223> OTHER INFORMATION: X = beta alanine

<400> SEQUENCE: 583

Arg Xaa Arg Arg Xaa Arg Arg Xaa Arg Arg Xaa Xaa Xaa  
 1 5 10

US 2010/0130591 A1

May 27, 2010

145

-continued

---

<210> SEQ ID NO 584  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 584  
ccactcagag ctcagatctt ctaacttcc 29  
  
<210> SEQ ID NO 585  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 585  
gggatccagt atacttacag gctcc 25  
  
<210> SEQ ID NO 586  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 586  
cttcactca gagctcagat cttctaa 27  
  
<210> SEQ ID NO 587  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 587  
acatcaagga agatggcatt tctagtttgg 30  
  
<210> SEQ ID NO 588  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 588  
ctccaacatc aaggaagatg gcatttctag 30  
  
<210> SEQ ID NO 589  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 589

US 2010/0130591 A1

May 27, 2010

146

-continued

---

ttctgtccaa gcccggttga aatc 24

<210> SEQ ID NO 590  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 590

caccacccat caccctcygt g 21

<210> SEQ ID NO 591  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 591

atcatctcgt tgatctctc aa 22

<210> SEQ ID NO 592  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 592

acatcaagga agatggcatt tctag 25

<210> SEQ ID NO 593  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 593

accagagtaa cagtctgagt aggagc 26

<210> SEQ ID NO 594  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 594

tcaaggaaga tggcatttct 20

<210> SEQ ID NO 595  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

147

-continued

---

<400> SEQUENCE: 595

cctctgtgat ttataaactt gat 23

<210> SEQ ID NO 596

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 596

atcatttttt ctcatacctt ctgct 25

<210> SEQ ID NO 597

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 597

ctcatacctt ctgcttgatg atc 23

<210> SEQ ID NO 598

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 598

tggcatttct agtttg 17

<210> SEQ ID NO 599

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 599

ccagagcagg tacctccaac atc 23

<210> SEQ ID NO 600

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 600

tgttcagctt ctgttagcca ctga 24

<210> SEQ ID NO 601

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 2010/0130591 A1

May 27, 2010

148

-continued

---

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 601

tttgtgtctt tctgagaaac 20

<210> SEQ ID NO 602  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 602

cgccgccatt tctcaacag 19

<210> SEQ ID NO 603  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 603

atctgtcaaa tcgcctgcag 20

<210> SEQ ID NO 604  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 604

tgtttttgag gattgctgaa 20

<210> SEQ ID NO 605  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 605

gctgaattat ttcttcccc 19

<210> SEQ ID NO 606  
<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 606

gcccaatgcc atcctgg 17

<210> SEQ ID NO 607  
<211> LENGTH: 26  
<212> TYPE: DNA



US 2010/0130591 A1

May 27, 2010

149

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 607  
  
ccaatgccat cctggagttc ctgtaa 26  
  
<210> SEQ ID NO 608  
<211> LENGTH: 31  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 608  
  
cattcaactg ttgcctccgg ttctgaaggt g 31  
  
<210> SEQ ID NO 609  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 609  
  
ctgaaggtgt tcttgtactt catcc 25  
  
<210> SEQ ID NO 610  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 610  
  
ctgttgccctc cggttctg 18  
  
<210> SEQ ID NO 611  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 611  
  
attctttcaa ctagaataaa ag 22  
  
<210> SEQ ID NO 612  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 612  
  
gccatcctgg agttcctgta agatacaaaa 30  
  
<210> SEQ ID NO 613

US 2010/0130591 A1

May 27, 2010

150

-continued

---

<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 613  
  
ccaatgccat cctggagttc ctgtaagata 30  
  
<210> SEQ ID NO 614  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 614  
  
gccgctgccc aatgccatcc tggagttcct 30  
  
<210> SEQ ID NO 615  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 615  
  
gtttgccgct gcccaatgcc atcctggagt 30  
  
<210> SEQ ID NO 616  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 616  
  
caacagtttg ccgctgcca atgccatcct 30  
  
<210> SEQ ID NO 617  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 617  
  
ctgacaacag tttgccgctg cccaatgcca 30  
  
<210> SEQ ID NO 618  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin  
  
<400> SEQUENCE: 618  
  
tgttctgaca acagtttgcc gctgccaat 30

US 2010/0130591 A1

May 27, 2010

151

-continued

---

```
<210> SEQ ID NO 619
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 619

caatgttctg acaacagttt gccgctgccc                               30

<210> SEQ ID NO 620
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 620

cattcaatgt tctgacaaca gtttgccgct                               30

<210> SEQ ID NO 621
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 621

tattttcttc ccagttgcat tcaatgttct                               30

<210> SEQ ID NO 622
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 622

gctgaattat ttcttcccca gttgcattca                               30

<210> SEQ ID NO 623
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 623

ggattgctga attatttctt ccccagttgc                               30

<210> SEQ ID NO 624
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense sequence to target splice site of
        proprocessed human dystrophin

<400> SEQUENCE: 624
```

US 2010/0130591 A1

May 27, 2010

152

-continued

---

tttgaggatt gctgaattat ttcttcccca 30

<210> SEQ ID NO 625  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 625

gtacttcac cccactgattc tgaattcttt 30

<210> SEQ ID NO 626  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 626

tcttgactt catccactg attctgaatt 30

<210> SEQ ID NO 627  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 627

tgttcttgta cttcatccca ctgattctga 30

<210> SEQ ID NO 628  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 628

cggttctgaa ggtgttcttg tacttcaccc 30

<210> SEQ ID NO 629  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

<400> SEQUENCE: 629

ctccggttct gaagggtgttc ttgtacttca 30

<210> SEQ ID NO 630  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Antisense sequence to target splice site of  
          proprocessed human dystrophin

US 2010/0130591 A1

May 27, 2010

153

-continued

---

<400> SEQUENCE: 630

tgccctccggt tctgaagggtg ttcttggtact

30

<210> SEQ ID NO 631

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 631

tgttgccctcc ggttctgaag gtgttctgt

30

<210> SEQ ID NO 632

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 632

aactgttgcc tccggttctg aagggtgttct

30

<210> SEQ ID NO 633

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 633

ttcaactgtt gcctccggtt ctgaagggt

30

<210> SEQ ID NO 634

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 634

taaagctctg gaaacctgaa aggaa

25

<210> SEQ ID NO 635

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense sequence to target splice site of  
proprocessed human dystrophin

<400> SEQUENCE: 635

ttcagcttct gtaagccagg caaga

25

<210> SEQ ID NO 636

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Antisense oligomer

US 2010/0130591 A1

May 27, 2010

154

-continued

<400> SEQUENCE: 636

ggccaaacct cggttacct gaaat

25

<210> SEQ ID NO 637

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: arginine-rich peptide transport

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: 1, 3, 4, 7, 9, 10

<223> OTHER INFORMATION: Xaa = K, R or arginine analog

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 2, 8

<223> OTHER INFORMATION: Xaa = a neutral amino acid C(=O) (CHR)n NH-,  
where n is 2 to 7 and each R is independently H or  
methyl

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: 5, 6, 11, 12

<223> OTHER INFORMATION: Xaa = an alpha-amino acid having a neutral  
aralkyl side chain

<400> SEQUENCE: 637

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
1 5 10

<210> SEQ ID NO 638

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: arginine-rich peptide transport

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 2, 6, 8, 12

<223> OTHER INFORMATION: Xaa = a neutral amino acid C(=O) (CHR)n NH-,  
where n is 2 to 7 and each R is independently H or  
methyl

<400> SEQUENCE: 638

Arg Xaa Arg Arg Arg Xaa Arg Xaa Arg Arg Arg Xaa  
1 5 10

<210> SEQ ID NO 639

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: arginine-rich peptide transport

<220> FEATURE:

<221> NAME/KEY: MOD\_RES

<222> LOCATION: 3, 5, 9

<223> OTHER INFORMATION: Xaa = a neutral amino acid C(=O) (CHR)n NH-,  
where n is 2 to 7 and each R is independently H or  
methyl

<400> SEQUENCE: 639

Arg Arg Xaa Arg Xaa Arg Arg Arg Xaa  
1 5

<210> SEQ ID NO 640

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

US 2010/0130591 A1

May 27, 2010

155

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 640

ccagagcttt acctgagaaa caag 24

<210> SEQ ID NO 641  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 641

ccagccactc agccagtgaa g 21

<210> SEQ ID NO 642  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 642

cgatccgtaa tgattgttct agcc 24

<210> SEQ ID NO 643  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 643

catttcattc aactgttgcc tccg 24

<210> SEQ ID NO 644  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 644

caatgctcct gacctctgtg c 21

<210> SEQ ID NO 645  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 645

gtctacaaca aagctcaggt cg 22

<210> SEQ ID NO 646  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 646

US 2010/0130591 A1

May 27, 2010

156

-continued

gcaatgttat ctgcttcctc caacc

25

<210> SEQ ID NO 647  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 647

gctcttttcc aggttcaagt gg

22

<210> SEQ ID NO 648  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 648

cttgacaga acttaccgac tgg

23

<210> SEQ ID NO 649  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 649

gcaggatttg gaacagaggc g

21

<210> SEQ ID NO 650  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 650

catctacatt tgtctgccac tgg

23

<210> SEQ ID NO 651  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 651

gtttcttcca aagcagcctc tcg

23

It is claimed:

1. A composition for use in producing skipping of exon 44 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 1-20, and capable of forming

with the complementary mRNA sequence in the dystrophin-gene exon 44, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

2. The composition of claim 1, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 8, 11 and 12.

3. The composition of claim 1, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.



US 2010/0130591 A1

May 27, 2010

157

4. A composition for use in producing skipping of exon 45 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 21-76 and 612 to 624, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 45, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

5. The composition of claim 4, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 27, 29, 34 and 39.

6. The composition of claim 5, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 29 and 34.

7. The composition of claim 4, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

8. A composition for use in producing skipping of exon 46 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 77-125, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 46, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

9. The composition of claim 8, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 77-105.

10. The composition of claim 9, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 82, 84-87, 90, 96, 98, 99 and 101.

11. The composition of claim 8, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

12. A composition for use in producing skipping of exon 47 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 126-169, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 47, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

13. The composition of claim 12, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 126-149.

14. The composition of claim 13, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 126, 128-130, 132, 144 and 146-149.

15. The composition of claim 12, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

16. A composition for use in producing skipping of exon 48 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 170-224 and 634, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 48, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

17. The composition of claim 16, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 170-201 and 634.

18. The composition of claim 17, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 176, 178, 181-183, 194 and 198-201.

19. The composition of claim 16, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

20. A composition for use in producing skipping of exon 49 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 225-266, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 49, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

21. The composition of claim 20, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 225-248.

22. The composition of claim 21, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 227, 229, 234, 236, 237 and 244-248.

23. The composition of claim 20, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

24. A composition for use in producing skipping of exon 50 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 267-308, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 50, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

25. The composition of claim 24, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 277, 287, 290 and 291.

US 2010/0130591 A1

May 27, 2010

158

26. The composition of claim 25, wherein the compound contains the sequence consisting of SEQ ID NO: 287.

27. The composition of claim 24, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

28. A composition for use in producing skipping of exon 51 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 309-371, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 51, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

29. The composition of claim 28, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 324, 326 and 327.

30. The composition of claim 29, wherein the compound contains the sequence consisting of SEQ ID NO: 327.

31. The composition of claim 28, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

32. A composition for use in producing skipping of exon 52 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 372-415, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 52, a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

33. The composition of claim 32, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 372-397.

34. The composition of claim 33, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 379-382, 384, 390 and 392-395.

35. The composition of claim 32, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

36. A composition for use in producing skipping of exon 53 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 416-475 and 625-633, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 53 a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

37. The composition of claim 36, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 428, 429 and 431.

38. The composition of claim 37, wherein the compound contains a sequence consisting of SEQ ID NO: 429.

39. The composition of claim 36, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

40. A composition for use in producing skipping of exon 54 in the processing of human dystrophin pre-processed mRNA, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 476-519, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 54 a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

41. The composition of claim 40, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 476-499.

42. The composition of claim 41, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 479-482, 484, 489 and 491-493.

43. The composition of claim 40, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

44. A composition for use in producing skipping of exon 55 in the processing of human dystrophin-gene pre-processed mRNA transcript, comprising

a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 520-569 and 635, and capable of forming with the complementary mRNA sequence in the dystrophin-gene exon 55 a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C.

45. The composition of claim 44, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 520-546 and 635.

46. The composition of claim 45, wherein the compound contains a sequence selected from the group consisting of SEQ ID NOS: 524-528, 537, 539, 540, 542 and 544.

47. The composition of claim 44, wherein the compound is conjugated to an arginine-rich peptide having a sequence selected from the group consisting of SEQ ID NOS: 570-578.

48. A method of treating muscular dystrophy in a subject, comprising administering to the subject an effective amount of a substantially uncharged antisense compound containing 20-35 morpholino subunits linked by phosphorus-containing intersubunit linkages joining a morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, comprising a sequence selected from the group consisting SEQ ID NOS: 1 to 569 and 612 to 635, and capable of forming with the complementary mRNA sequence in a dystrophin-gene exon a heteroduplex structure between said compound and mRNA having a T<sub>m</sub> of at least 45° C., wherein the exon is selected from the group consisting of exons 44-55.

49. The method of claim 48, wherein the muscular dystrophy is Duchenne's muscular dystrophy (DMD).

US 2010/0130591 A1

May 27, 2010

159

**50.** The method of claim **48**, wherein the muscular dystrophy is Becker muscular dystrophy (BMD).

**51.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 1-20, and the exon is exon 44.

**52.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 21-76 and 612 to 624, and the exon is exon 45.

**53.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 77-125, and the exon is exon 46.

**54.** The method of claim **48**, wherein the sequence selected from the group consisting SEQ ID NOS: 126-169, and the exon is exon 47.

**55.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 170-224 and 634, and the exon is exon 48.

**56.** The method of claim **48**, wherein the sequence selected from the group consisting SEQ ID NOS: 225-266, and the exon is exon 49.

**57.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 267-308, and the exon is exon 50.

**58.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 309-371, and the exon is exon 51.

**59.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 372-415, and the exon is exon 52.

**60.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 416-475 and 625-633, and the exon is exon 53.

**61.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 476-519, and the exon is exon 54.

**62.** The method of claim **48**, wherein the sequence is selected from the group consisting SEQ ID NOS: 520-569 and 635, and the exon is exon 55.

**63.** The method of claim **48**, wherein the sequence comprises SEQ ID NO:287.

**64.** The method of claim **48**, wherein the compound is conjugated to an arginine-rich peptide.

**65.** The method of claim **64**, wherein the arginine-rich peptide comprises a sequence selected from the group consisting of SEQ ID NOS: 570-578.

\* \* \* \* \*